

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/90074 A2

(51) International Patent Classification⁵: C07D 213/74,
213/89, 217/22, 215/38, 209/08, 263/58, 235/30, 277/62,
239/42, A61K 31/33

(74) Agent: THALSØ-MADSEN, Birgit; Leo Pharmaceutical
Products, Industriparken 55, DK-2750 Ballerup (DK).

(21) International Application Number: PCT/DK01/00346

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(22) International Filing Date: 18 May 2001 (18.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/205,579 22 May 2000 (22.05.2000) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):
LEO PHARMACEUTICAL PRODUCTS LTD. A/S
(LØVENS KEMISKE FABRIK PRODUKTIONSAK-
TIESELSKAB) [DK/DK]; Industriparken 55, DK-2750
Ballerup (DK).

Published:

— *without international search report and to be republished
upon receipt of that report*

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): HORNEMAN, Anne,
Marie [DK/DK]; Ida Tesdorfsvej 4B, 1.tv, DK-3050
Humblebæk (DK).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 01/90074 A2

(54) Title: BENZOPHENONES AS INHIBITORS OF IL-1 β AND TNF- α

(57) Abstract: Heteroaryl aminobenzophenones of general formula (I) inhibit interleukin-1 β and TNF- α and may therefore be use-
ful in the therapy of inflammatory diseases and conditions.

BENZOPHENONES AS INHIBITORS OF IL-1 β AND TNF- α

FIELD OF THE INVENTION

- 5 The present invention relates to a novel class of compounds, including heteroaryl aminobenzophenone derivatives, which show anti-inflammatory effects, to pharmaceutical compositions containing these compounds, and to their use in the treatment and prophylaxis of inflammatory diseases.

10 BACKGROUND OF THE INVENTION

- A series of aminobenzophenones (e.g. 4-(2-amino-4-nitrophenylamino)benzophenone) have been described previously (Hussein, F.A. et al., Iraqi J. Sci., 22, 54-66 (1981)). In this publication, however, there is no description of any potential therapeutic use of such compounds. WO 98/32730 discloses aminobenzophenone inhibitors of interleukin 1 β (IL-1 β) and tumour necrosis factor α (TNF- α) secretion *in vitro*, and indicates the potential utility of these compounds in the treatment of inflammatory diseases in which the production of pro-inflammatory cytokines is involved in the pathogenesis, e.g. asthma, rheumatoid arthritis, psoriasis, contact dermatitis, and atopic dermatitis. Furthermore the compounds disclosed in WO 98/32730 were tested *in vivo* for anti-inflammatory properties in the 12-O-tetradecanoylphorbol-13-acetate (TPA) induced murine chronic skin inflammation model (De Young, L.M. et al., *Agents Actions* 26, 335-341 (1989); Carlson, R.P. et al., *Agents Actions* 17, 197-204 (1985); Alford, J.G. et al., *Agents Action* 37, (1992); Stanley, P.L. et al., *Skin Pharmacol.* 4, 262-271 (1991)). In this chronic skin inflammation model the compounds had the same potency compared to the reference compound hydrocortisone.


It is the object of the present invention is to provide further pharmacologically active benzophenone derivatives which differ structurally from those disclosed in WO 98/32730.

30 SUMMARY OF THE INVENTION

It has surprisingly been found that novel benzophenone derivatives are potent inhibitors of Interleukin 1 β (IL-1 β) and tumour necrosis factor α (TNF- α) secretion *in vitro*, making them potentially useful for the treatment and/or prevention of inflammatory diseases and other conditions in which the secretion and regulation of cytokines or more specifically

interleukin 1 β (IL-1 β) and tumour necrosis factor α (TNF- α) are involved in the pathogenesis. The inhibition or downregulation of the cytokines is possibly due to an inhibition of MAP kinases, more specifically the p38 MAP kinase, a stress-activated protein which is an important element in the signal transduction pathway leading to the production of pro-inflammatory cytokines.

Accordingly, the present invention relates to a compound with the general formula I



(I)

- wherein R₁ is selected from the group consisting of halogen, haloalkyl, hydroxy, hydroxyalkyl, hydroxyalkyloxy, mercapto, cyano, carboxy, nitro, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, aryl, heteroaryl, aralkyl, alkylaryl, alkoxy, aralkoxy, alkylthio, alkoxycarbonyl, alkylcarbonyloxy, alkoxycarbonyloxy, alkylsulfonyloxy, alkylloxysulfonyl, alkylcarbonylamino, aminocarboaminoalkyl, aminosulfonyl, alkylsulfonylamino, alkanoyl, alkylcarbonyl, -NR₉R₁₀ or -CONR₉R₁₀, wherein R₉ and R₁₀ are the same or different and individually represent hydrogen, alkyl or aryl;

- R₂ represents one or more, same or different substituents selected from the group consisting of hydrogen, halogen, haloalkyl, hydroxy, hydroxyalkyl, hydroxyalkyloxy, mercapto, cyano, carboxy, nitro, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, aryl, heteroaryl, aralkyl, alkylaryl, alkoxy, aralkoxy, alkylthio, alkoxycarbonyl, alkylcarbonyloxy, alkoxycarbonyloxy, alkylsulfonyloxy, alkylloxysulfonyl, alkylcarbonylamino, aminocarboaminoalkyl, aminosulfonyl, alkylsulfonylamino, alkanoyl, alkylcarbonyl, -NR₉R₁₀ or -CONR₉R₁₀, wherein R₉ and R₁₀ are the same or different and individually represent hydrogen, alkyl or aryl;

- R₃ represents one or more, same or different substituents selected from the group consisting of hydrogen, halogen, haloalkyl, hydroxy, hydroxyalkyl, mercapto, cyano, nitro, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, aralkyl, alkylaryl, alkoxy, aralkoxy, alkylthio, alkoxycarbonyl, alkylcarbonylamino, alkylcarbonyloxy, alkoxycarbonyloxy, alkylcarbonyl -

NR₉R₁₀ or -CONR₉R₁₀, wherein R₉ and R₁₀ are the same or different and individually represent hydrogen, alkyl or aryl;

R₄ represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, carboxy or aryl;

5

R₅ represents a heteroaromatic mono- or bicyclic ring system comprising 1-4 heteroatoms, except for triazine, said ring system being optionally substituted by hydrogen, halogen, haloalkyl, hydroxy, hydroxyalkyl, hydroxyalkyloxy, mercapto, cyano, carboxy, nitro, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, aryl, heteroaryl, aralkyl, alkylaryl, alkoxy, aralkoxy, alkylthio, alkoxycarbonyl, alkylcarbonyloxy, alkoxy carbonyloxy, alkylsulfonyloxy, alkylloxysulfonyl, alkylcarbonylamino, aminocarboaminoalkyl, aminosulfonyl, alkylsulfonylamino, alkanoyl, alkylcarbonyl, -NR₉R₁₀ or -CONR₉R₁₀, wherein R₉ and R₁₀ are the same or different and individually represent hydrogen, alkyl or aryl;

15

X represents oxygen, sulphur, N-OH or NR₁₁ wherein R₁₁ is hydrogen or alkyl;

or pharmaceutically acceptable salts hydrates, solvates or esters thereof as well as N-oxides wherein an N-atom of R₅ is oxidised.

20

DETAILED DESCRIPTION OF THE INVENTION

Definitions

25 In the present context, the term "alkyl" is intended to indicate a univalent radical derived from straight or branched alkane by removing a hydrogen atom from any carbon atom. The alkyl chain typically comprises 1-10 carbon atoms, in particular 1-6 carbon atoms. The term includes the subclasses normal alkyl (*n*-alkyl), secondary and tertiary alkyl, such as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *sec*.-butyl, *tert*.-butyl, pentyl, isopentyl, hexyl and isohexyl.

30

The term "haloalkyl" is intended to indicate an alkyl radical as defined above substituted by one or more halogens such as chloro, fluoro, bromo or iodo.

35 The term "hydroxyalkyl" is intended to indicate an alkyl radical as defined above substituted by one or more hydroxy groups.

The term "alkoxy" is intended to indicate a radical of formula OR', wherein R' is alkyl as

defined above, e.g. methoxy, ethoxy, propoxy, butoxy, etc.

The term "hydroxyalkyloxy" is intended to indicate an alkoxy group as defined above substituted by one or more hydroxy groups.

5

The term "alkenyl" is intended to indicate a mono-, di-, tri-, tetra- or pentaunsaturated alkane radical typically comprising 2-10 carbon atoms, in particular 2-6 carbon atoms, e.g. ethenyl, propenyl, butenyl, pentenyl or hexenyl. The term "alkynyl" is intended to indicate an alkane radical comprising 1-5 triple C-C bonds, the alkane chain typically comprising 2-10 carbon atoms, in particular 2-6 carbon atoms, such as ethynyl, propynyl, butynyl, pentynyl or hexynyl.

10

The term "alkoxycarbonyl" is intended to indicate a radical of formula -COOR' wherein R' is alkyl as defined above, e.g. methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, isopropoxycarbonyl, etc.

15

The term "cycloalkyl" is intended to indicate a saturated cycloalkane radical typically comprising 3-10 carbon atoms, in particular 3-8 carbon atoms, e.g. cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl. The term "cycloalkenyl" is intended to indicate mono-, di- tri- or tetraunsaturated cycloalkane radicals, e.g. cyclopropenyl, cyclobutenyl, cyclopentenyl or cyclohexenyl. The term "heterocycloalkyl" is intended to indicate a cycloalkane radical as defined above, comprising one or more heteroatoms selected from O, N, or S.

20

The term "aryl" is intended to include radicals of carbocyclic aromatic rings, in particular 5- or 6-membered rings, optionally fused bicyclic rings, e.g. phenyl or naphthyl. The term "heteroaryl" is intended to include radicals of heterocyclic aromatic rings, in particular 5- or 6-membered rings with 1-4 heteroatoms selected from O, S and N, or optionally fused bicyclic rings with 1-4 heteroatoms, e.g. pyridyl, quinolyl, isoquinolyl, indolyl, tetrazolyl, thiazolyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thienyl, pyrazinyl, isothiazolyl, benzimidazolyl and benzofuranyl.

25

30

The term "alkylcarbonyloxy" refers to a radical of formula R'-COO- , wherein R' is alkyl as indicated above. The term "alkoxycarbonyloxy" refers to a radical of formula R'O-COO- , wherein R' is alkyl as defined above. The term "alkylsulfonyloxy" refers to a radical of formula $\text{R'-(SO}_2\text{)-O-}$, wherein R' is alkyl as defined above. The term "alkyloxysulfonyl" refers to a radical of formula $\text{R'O-(SO}_2\text{)-}$, wherein R' is alkyl as defined above.

35

The term "aralkyl" is intended to indicate an aromatic ring with an alkyl side chain as defined above, e.g. benzyl. The term "alkylaryl" is intended to indicate an alkyl radical as defined above comprising an aromatic side chain.

5

The term "halogen" is intended to indicate fluoro, chloro, bromo or iodo.

The term "alkylthio" is intended to indicate a radical of the formula -SR, where R is alkyl as defined above and includes methylthio, ethylthio, n-propylthio, and 2-propylthio.

- 10 The term "alkylamino" is intended to indicate a radical of the formula -NHR or -NR₂, where R is alkyl as defined above having from 1-6 carbon atoms and includes, for example, methylamino, dimethylamino, di-(n-propyl)amino, and n-butyl(ethyl)amino.

The term "carbamoyl" is intended to indicate the group -CONH₂, -CONHR, and -

- 15 OCONRR' where R and R' represent alkyl as defined above.

The term "pharmaceutically acceptable salt" is intended to indicate alkali metal or alkaline earth metal salts, for instance sodium, potassium, magnesium or calcium salts, as well as silver salts and salts with suitable organic or inorganic acids such as hydrochloric, hydrobromic, hydroiodic, sulfuric, nitric, phosphoric, acetic, lactic, maleic, phthalic, citric, propionic, benzoic, glutaric, gluconic, methanesulfonic, salicylic, succinic, tartaric, toluenesulfonic, sulfamic or fumaric acid.

- 25 The term "pharmaceutically acceptable esters" is intended to indicate easily hydrolysable esters such as alkanoyloxyalkyl, aralkanoyloxyalkyl, aroyloxyalkyl, e.g. acetoxymethyl, pivaloyloxymethyl, benzoyloxymethyl esters and the corresponding 1'-oxyethyl derivatives, or alkoxycarbonyloxyalkyl esters, e.g. methoxycarbonyloxymethyl esters and ethoxycarbonyloxymethyl esters and the corresponding 1'-oxyethyl derivatives, or lactonyl esters, e.g. phthalidyl esters, or dialkylaminoalkyl esters, e.g. dimethylaminoethyl esters.
- 30 Easily hydrolysable esters include *in vivo* hydrolysable esters of the compounds of formula I. Such esters may be prepared by conventional methods known to persons skilled in the art, such as method disclosed in GB patent No. 1 490 852 incorporated herein by reference.

- 35 "p38 MAP kinase" is a stress-activated protein kinase existing in several isoforms (p38 α , p38 β , p38 β 2, p38 γ and p38 δ). The p38 MAP kinase is activated by different stimuli including heat, chemical, osmotic, pH and oxidative stress, growth factor withdrawal, high or low glucose and ultraviolet radiation. p38 is also stimulated by agents that mediate the

initial physiological response to Injury, infection and inflammation, such as LPS and pro-inflammatory cytokines IL-1 β , TNF- α , FasL, CD40L and TGF- β . Like other MAP kinases, p38 is phosphorylated by kinases, including MKK3, MEK6 and MKK6, on a threonine and tyrosine in an activation loop (Thr-Xaa-Tyr) close to the ATP and substrate binding site. In turn, p38 phosphorylates and activates the serine-threonine protein kinases MAPKAP kinase-2, MAPKAP kinase-3, MAPKAP kinase-5, MNK-1 and MSK-1. It has been established that activation of p38 regulates cytokine biosynthesis in many cell types either directly by phosphorylating and activating transcription factors involved in the expression of cytokines or indirectly, e.g. by phosphorylating MSK-1 which, when activated, activates the transcription factor CREB. It has also been shown that certain pyridinyl imidazoles, e.g. SB203580, inhibit the production of IL-1 β and TNF- α from LPS-treated human monocytes, are inhibitors of p38 kinase. It has therefore been concluded that p38 constitutes a potentially highly interesting target for the development of anti-inflammatory compounds (cf. JC Lee et al., *Immunopharmacology* 47, 2000, pp. 185-201 and references reviewed therein; PR Young, "Specific Inhibitors of p38 MAP kinase" In *Signaling Networks and Cell Cycle Control: The Molecular Basis of Cancer and Other Diseases*, JS Gutkind (Ed.), Humana Press, Inc., Totowa, NJ, and references reviewed therein).

Preferred embodiments of the invention

In compounds of formula I,

R₁ preferably represents a substituent selected from the group consisting of halogen, hydroxy, mercapto, trifluoromethyl, amino, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, (C₁-C₆)alkylamino, (C₁-C₆)alkoxycarbonyl, cyano, -CONH₂, phenyl, and nitro, in particular fluoro, chloro, bromo, hydroxy, trifluoromethyl, amino, (C₁-C₃)alkyl, (C₂-C₃)alkenyl, (C₁-C₃)alkoxy, (C₁-C₃)alkoxycarbonyl, cyano, or -CONH₂;

R₂ represents one or more, same or different substituents selected from the group consisting of hydrogen, halogen, hydroxy, mercapto, trifluoromethyl, amino, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, (C₁-C₆)alkylamino, (C₁-C₆)alkoxycarbonyl, cyano, -CONH₂, phenyl, and nitro; in particular hydrogen, fluoro, chloro, bromo, hydroxy, trifluoromethyl, amino, (C₁-C₃)alkyl, (C₂-C₃)alkenyl, (C₁-C₃)alkoxy, (C₁-C₃)alkoxycarbonyl, cyano, or -CONH₂;

R_3 represents one or more, same or different substituents selected from the group consisting of hydrogen, halogen, hydroxy, mercapto, trifluoromethyl, amino, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, (C_1-C_6) alkoxy, (C_1-C_6) alkylthio, (C_1-C_6) alkylamino, (C_1-C_6) alkoxycarbonyl, cyano, $-CONH_2$, phenyl, and nitro, in particular hydrogen, fluoro, chloro, bromo, hydroxy, trifluoromethyl, amino, (C_1-C_3) alkyl, (C_2-C_3) alkenyl, and (C_1-C_3) alkoxy;



R_4 represents hydrogen, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, or (C_3-C_6) cycloalkyl or cycloalkenyl, in particular hydrogen, (C_1-C_4) alkyl, or (C_2-C_4) alkenyl;

X represents oxygen or NH;


In the mono- or bicyclic heteroaromatic ring system represented by R_5 , the heteroatom(s) may be selected from N, S or O. Each ring preferably comprises 5 or 6 ring atoms.

Examples of suitable heteroaromatic ring systems are selected from the group consisting of pyridyl, pyrazinyl, pyrimidinyl, quinolyl, isoquinolyl, quinazolinyl, pyridazinyl, phthalazinyl, purinyl, quinoxalyl, allopurinyl, benzofuranyl, isobenzofuranyl, benzimidazolyl, benzoxazolyl, benzothienyl, isobenzothienyl, benzothiazolyl, benzoisothiazolyl, benzotriazolyl, indolizynyl, isoindolyl, indolyl, indazolyl, triazolyl, oxazolyl, thiadiazolyl, thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl or isoxazolyl.



R_5 is preferably selected from heteroaromatic ring systems comprising 1 or 2 nitrogen atoms, e.g. rings of the structures:





The R_5 ring-systems may optionally be oxidised to the corresponding N-oxides as exemplified below for the isoquinoline ring system:



5




10



- 15 R_6 and R_7 represent one or more, same or different substituents selected from the group consisting of hydrogen, halogen, hydroxy, mercapto, trifluoromethyl, cyano, carboxy, carbamoyl, amino, nitro, (C_1-C_{10}) alkyl, (C_2-C_{10}) alkenyl, (C_3-C_8) cycloalkyl or-cycloalkenyl, (C_1-C_{10}) alkoxy, (C_1-C_{10}) alkylthio, (C_1-C_{10}) alkoxycarbonyl, and phenyl. In particular R_6 and R_7 represent one or more, same or different substituents selected from
- 20 the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, trifluoromethyl, amino, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, (C_1-C_6) alkoxy, (C_1-C_6) alkoxycarbonyl, cyano, carboxy, and $-CONH_2$; and

25



R_8 represents hydrogen, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, or (C_3-C_6) cycloalkyl or -cycloalkenyl.

- 30 In compounds of formula I it is more preferred that

R_1 represents a substituent selected from the group consisting of fluoro, chloro, bromo, hydroxy, methyl, and methoxy;

R₂ represents one or more, same or different substituents selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, trifluoromethyl, methyl, ethyl, and methoxy;

- 5 R₃ represents one or more, same or different substituents selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, methyl, and methoxy;

R₄ represents hydrogen, methyl, or ethyl;

- 10 R₅ is selected from the group consisting of substituted or non-substituted 3-pyridyl, 2-pyridyl, 3-quinolyl, 4-isoquinolyl, 4-indolyl, 5-indolyl, 6-indolyl and 7-indolyl moieties, and the corresponding N-oxides;

- R₆ and R₇ represent one or more, same or different substituents selected from the group
15 consisting of hydrogen, fluoro, chloro, bromo, hydroxy, methyl, methoxy, cyano, and carboxy;

R₈ represents hydrogen, (C₁-C₄)alkyl, (C₂-C₆)alkenyl, such as allyl; and more preferably, R₈ represents hydrogen, methyl, ethyl, allyl, propyl, or t-butyl; and/or

20

X represents oxygen.

The phenyl group of R₁, R₂, R₃, R₆ and R₇ may optionally be substituted, e.g. with hydroxy, amino, nitro, cyano or halogen, preferably fluoro or chloro.

25

Specific compounds of the Invention are:

- 2-Chloro-2'-methyl-4-(4-pyridylamino)benzophenone (Compound 101),
2-Chloro-2'-methyl-4-(2-pyridylamino)benzophenone (Compound 102),
30 2-Chloro-2'-methyl-4-(5-nitro-2-pyridylamino)benzophenone (Compound 103),
4-(6-Amino-5-nitro-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 104),
6-Chloro-2-(3-chloro-4-(2-methylbenzoyl)phenylamino) isonicotinic acid (Compound 105),
4-(6-carbonitrile-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 106),
2-Chloro-2'-methyl-4-(3-pyridylamino)benzophenone (Compound 107),
35 2-Chloro-4-(5,6-diamino-2-pyridylamino)-2'-methylbenzophenone (Compound 108),
2-Chloro-2'-methyl-4-(3-nitro-2-pyridylamino)benzophenone (Compound 109),
4-(3-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 110),

- 4-(5-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 111),
2-Chloro-4-(4-isoquinolylamino)-2'-methylbenzophenone (Compound 114),
t-Butyl 5-(3-chloro-4-(2-methylbenzoyl)phenylamino) nicotinoate (Compound 115),
2-Chloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 116),
5 2-Chloro-2'-methyl-4-(6-methyl-3-pyridylamino)benzophenone (Compound 117),
4-(5-Bromo-3-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 118),
4-(5-Carbonitrile-3-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 119),
4-(3-Bromo-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 120),
2-Chloro-2'-methyl-4-(8-quinolylamino)benzophenone (Compound 121),
10 2-Chloro-4-(6-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 122),
4-(4-Bromo-1-isoquinolylamino)-2-chloro-2'-methylbenzophenone (Compound 123),
2-Chloro-2'-methyl-4-(2-methyl-5-trifluoromethyl-3-pyridylamino)benzophenone
(Compound 124),
2-Chloro-2'-methyl-4-(3-quinolylamino)benzophenone (Compound 125),
15 2-Chloro-4-(4-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 126),
2-Chloro-4-(2-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 127),
2-Chloro-2'-methyl-4-(2-methyl-6-quinolylamino)benzophenone (Compound 129),
2-Chloro-4-(7-chloro-4-quinolylamino)-2'-methylbenzophenone (Compound 130),
2-Chloro-2'-methyl-4-(2-quinolylamino)benzophenone (Compound 131),
20 2-Chloro-2'-methyl-4-(4-quinolylamino)benzophenone (Compound 132),
2-Chloro-2'-methyl-4-(1-methyl-7-indolylamino)benzophenone (Compound 133),
2-Chloro-2'-methyl-4-(1-methyl-5-indolylamino)benzophenone (Compound 134),
2-Chloro-2',5'-dimethyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 135),
2-Chloro-2',5'-dimethyl-4-(4-isoquinolylamino)benzophenone (Compound 136),
25 2-Chloro-4-(4-isoquinolylamino)-2',4',5'-trimethylbenzophenone (Compound 137),
2,3'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 138),
2-Fluoro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 139),
2,4'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 140),
2-Chloro-4'-fluoro-4-(4-isoquinolylamino)-2'-methylbenzophenone (Compound 141),
30 4'-n-Butyl-2-chloro-4-(4-isoquinolylamino)-2'-methylbenzophenone (Compound 142),
2-Chloro-4-(5-isoquinolylamino)-2'-methylbenzophenone (Compound 143),
2-Chloro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone (Compound 144),
2-Fluoro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone (Compound 145),
2,4'-Dichloro-2'-methyl-4-(1-methyl-7-indolylamino)benzophenone (Compound 146),
35 2-Chloro-4-(1-methyl-7-indolylamino)-2',4',5'-trimethylbenzophenone (Compound 147),
2-Chloro-2',5'-dimethyl-4-(1-methyl-7-indolylamino)benzophenone (Compound 148),
2-Chloro-4-(3-ethoxy-4-isoquinolylamino)-2'-methylbenzophenone (Compound 149),
2-Chloro-4-(1-ethoxy-4-isoquinolylamino)-2'-methylbenzophenone (Compound 150),

- 4-(2-Benzoxazolylamino)-2-chloro-2'-methylbenzophenone (Compound 151),
4-(1-Methyl-2-benzimidazolylamino)-2-chloro-2'-methylbenzophenone (Compound 152),
4-(2-Benzothiazolylamino)-2-chloro-2'-methylbenzophenone (Compound 153),
2-Chloro-2'-methyl-4-(2-pyrimidylamino)benzophenone (Compound 154),
5 2-Chloro-2'-methyl-4-(7-methyl-purin-6-ylamino)benzophenone (Compound 155),
2-Chloro-2'-methyl-4-(2-methyl-5-benzothiazolylamino)benzophenone (Compound 156),
2-Chloro-2'-methyl-4-(pyrazin-2-ylamino)benzophenone (Compound 157),
2-Chloro-2'-methyl-4-(5-pyrimidylamino)benzophenone (Compound 158),
2-Chloro-2'-methyl-4-(5-nitro-2-thiazolylamino)benzophenone (Compound 159),
10 2-Chloro-2'-methyl-4-((4-methyl-3-nitro-(1,2,4-triazol-5-ylamino)))benzophenone
(Compound 160)

- and the N-oxides thereof wherein the nitrogen atom of the heterocyclic R₅ substituent is
15 specifically oxidised.

More preferred compounds are:


- 2-Chloro-2'-methyl-4-(3-pyridylamino)benzophenone (Compound 107),
4-(3-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 110),
20 2-Chloro-4-(4-isoquinolylamino)-2'-methylbenzophenone (Compound 114),
2-Chloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 116),
2-Chloro-2'-methyl-4-(6-methyl-3-pyridylamino)benzophenone (Compound 117),
4-(5-Bromo-3-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 118),
2-Chloro-2'-methyl-4-(8-quinolylamino)benzophenone (Compound 121),
25 2-Chloro-4-(6-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 122),
2-Chloro-2'-methyl-4-(3-quinolylamino)benzophenone (Compound 125),
2-Chloro-4-(4-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 126),
2-Chloro-2'-methyl-4-(1-methyl-7-indolylamino)benzophenone (Compound 133),
2-Chloro-2'-methyl-4-((1-methyl-5-indolylamino)benzophenone (Compound 134),
30 2-Chloro-2',5'-dimethyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 135)
2-Chloro-2',5'-dimethyl-4-(4-isoquinolylamino)benzophenone (Compound 136),
2-Chloro-4-(4-isoquinolylamino)-2',4',5'-trimethylbenzophenone (Compound 137),
2-Fluoro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 139),
2,4'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 140),
35 2-Chloro-4'-fluoro-4-(4-isoquinolylamino)-2'-methylbenzophenone (Compound 141),
2-Chloro-4-(5-isoquinolylamino)-2'-methylbenzophenone (Compound 143),
2-Chloro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone (Compound 144),
2-Fluoro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone (Compound 145),

2-Chloro-4-((3-ethoxy)-4-isoquinolylamino)-2'-methylbenzophenone (Compound 149),
 2-Chloro-4-((1-ethoxy)-4-isoquinolylamino)-2'-methylbenzophenone (Compound 150)


and the N-oxides

- 5 2-Chloro-2'-methyl-4-((2-pyridyl-N-oxide)amino)benzophenone (Compound 112),
 2-Chloro-2'-methyl-4-((3-pyridyl-N-oxide)amino)benzophenone (Compound 113),
 2-Chloro-4-((4-isoquinolyl-N-oxide)amino)-2'-methylbenzophenone (Compound 128),

- Compounds of formula Ia wherein X=S and wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇ and R₈
 10 are as indicated above, and compounds of formula Ib wherein X=N-OH and wherein R₁,
 R₂, R₃, R₄, R₅, R₆, R₇ and R₈ are as indicated above are also generally preferred.



Ia



Ib

- 15 Specific compounds of formula Ia are:

- 2-Chloro-2'-methyl-4-(4-pyridylamino)thiobenzophenone,
 2-Chloro-2'-methyl-4-(2-pyridylamino)thiobenzophenone,
 2-Chloro-2'-methyl-4-(5-nitro-2-pyridylamino)thiobenzophenone,
 20 4-(6-Amino-5-nitro-2-pyridylamino)-2-chloro-2'-methyl(thiobenzophenone),
 6-Chloro-2-(3-chloro-4-(2-methylthiobenzoyl)phenylamino) isonicotinic acid,
 4-(6-carbonitrile-2-pyridylamino)-2-chloro-2'-methyl(thiobenzophenone),
 2-Chloro-2'-methyl-4-(3-pyridylamino)thiobenzophenone,
 2-Chloro-4-(5,6-diamino-2-pyridylamino)-2'-methyl(thiobenzophenone),
 25 2-Chloro-2'-methyl-4-(3-nitro-2-pyridylamino)thiobenzophenone,
 4-(3-Amino-2-pyridylamino)-2-chloro-2'-methyl(thiobenzophenone),
 4-(5-Amino-2-pyridylamino)-2-chloro-2'-methyl(thiobenzophenone),
 2-Chloro-4-(4-isoquinolylamino)-2'-methyl(thiobenzophenone),
 t-Butyl 5-(3-chloro-4-(2-methylthiobenzoyl)phenylamino)nicotinoate,
 30 2-Chloro-2'-methyl-4-(4-methyl-3-pyridylamino)thiobenzophenone,
 2-Chloro-2'-methyl-4-(6-methyl-3-pyridylamino)thiobenzophenone,

- 4-(5-Bromo-3-pyridylamino)-2-chloro-2'-methyl(thiobenzophenone),
4-(5-Carbonitrile-3-pyridylamino)-2-chloro-2'-methyl(thiobenzophenone),
4-(3-Bromo-2-pyridylamino)-2-chloro-2'-methyl(thiobenzophenone),
2-Chloro-2'-methyl-4-(8-quinolylamino)thiobenzophenone,
5 2-Chloro-4-(6-ethoxy-3-pyridylamino)-2'-methyl(thiobenzophenone),
4-(4-Bromo-1-isoquinolylamino)-2-chloro-2'-methyl(thiobenzophenone),
2-Chloro-2'-methyl-4-(2-methyl-5-trifluoromethyl-3-pyridylamino)thiobenzophenone,
2-Chloro-2'-methyl-4-(3-quinolylamino)thiobenzophenone,
2-Chloro-4-(4-ethoxy-3-pyridylamino)-2'-methyl(thiobenzophenone),
10 2-Chloro-4-(2-ethoxy-3-pyridylamino)-2'-methyl(thiobenzophenone),
2-Chloro-2'-methyl-4-(2-methyl-6-quinolylamino)thiobenzophenone,
2-Chloro-2'-methyl-4-(2-quinolylamino)thiobenzophenone,
2-Chloro-4-(7-chloro-4-quinolylamino)-2'-methyl(thiobenzophenone),
2-Chloro-2'-methyl-4-(4-quinolylamino)thiobenzophenone,
15 2-Chloro-2'-methyl-4-(1-methyl-7-indolylamino)thiobenzophenone, and
2-Chloro-2'-methyl-4-(1-methyl-5-indolylamino)thiobenzophenone,
2-Chloro-2',5'-dimethyl-4-(4-methyl-3-pyridylamino)thiobenzophenone,
2-Chloro-2',5'-dimethyl-4-(4-isoquinolylamino)thiobenzophenone,
2-Chloro-4-(4-isoquinolylamino)-2',4',5'-trimethyl(thiobenzophenone),
20 2,3'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)thiobenzophenone,
2-Fluoro-2'-methyl-4-(4-methyl-3-pyridylamino)thiobenzophenone,
2,4'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)thiobenzophenone,
2-Chloro-4'-fluoro-4-(4-isoquinolylamino)-2'-methyl(thiobenzophenone),
4'-*n*-Butyl-2-chloro-4-(4-isoquinolylamino)-2'-methyl(thiobenzophenone),
25 2-Chloro-4-(5-isoquinolylamino)-2'-methyl(thiobenzophenone),
2-Chloro-4-(4-isoquinolylamino)-4'-methoxy-2'-methyl(thiobenzophenone),
2-Fluoro-4-(4-isoquinolylamino)-4'-methoxy-2'-methyl(thiobenzophenone),
2,4'-Dichloro-2'-methyl-4-(1-methyl-7-indolylamino)thiobenzophenone,
2-Chloro-4-(1-methyl-7-indolylamino)-2',4',5'-trimethyl(thiobenzophenone),
30 2-Chloro-2',5'-dimethyl-4-(1-methyl-7-indolylamino)thiobenzophenone,
2-Chloro-4-(3-ethoxy-4-isoquinolylamino)-2'-methyl(thiobenzophenone),
2-Chloro-4-(1-ethoxy-4-isoquinolylamino)-2'-methyl(thiobenzophenone),
4-(2-Benzoxazolylamino)-2-chloro-2'-methyl(thiobenzophenone),
4-(1-Methyl-2-benzimidazolylamino)-2-chloro-2'-methyl(thiobenzophenone),
35 4-(2-Benzothiazolylamino)-2-chloro-2'-methyl(thiobenzophenone),
2-Chloro-2'-methyl-4-(2-pyrimidylamino)thiobenzophenone,
2-Chloro-2'-methyl-4-(2-methyl-purin-6-ylamino)thiobenzophenone,
2-Chloro-2'-methyl-4-(2-methyl-5-benzothiazolylamino)thiobenzophenone,

2-Chloro-2'-methyl-4-(pyrazin-2-ylamino)thiobenzophenone,
2-Chloro-2'-methyl-4-(5-pyrimidylamino)thiobenzophenone

and salts thereof with pharmaceutically acceptable acids, hydrates and solvates.

5


Specific compound of formula Ib are:

- 2-Chloro-2'-methyl-4-(4-pyridylamino)benzophenone oxime,
2-Chloro-2'-methyl-4-(2-pyridylamino)benzophenone oxime,
10 2-Chloro-2'-methyl-4-(5-nitro-2-pyridylamino)benzophenone oxime,
4-(6-Amino-5-nitro-2-pyridylamino)-2-chloro-2'-methylbenzophenone oxime,
6-Chloro-2-((3-chloro-4-((hydroxyimino)(2-methylphenyl)methyl))phenylamino)
isonicotinic acid,
4-(6-carbonitrile-2-pyridylamino)-2-chloro-2'-methylbenzophenone oxime,
15 2-Chloro-2'-methyl-4-(3-pyridylamino)benzophenone oxime,
2-Chloro-4-(5,6-diamino-2-pyridylamino)-2'-methylbenzophenone oxime,
2-Chloro-2'-methyl-4-(3-nitro-2-pyridylamino)benzophenone oxime,
4-(3-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone oxime,
4-(5-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone oxime,
20 2-Chloro-4-(4-isoquinolylamino)-2'-methylbenzophenone oxime,
t-Butyl 5-((3-chloro-4-((hydroxyimino)(2-methylphenyl)methyl))phenylamino)
nicotinoate,
2-Chloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone oxime,
2-Chloro-2'-methyl-4-(6-methyl-3-pyridylamino)benzophenone oxime,
25 4-(5-Bromo-3-pyridylamino)-2-chloro-2'-methylbenzophenone oxime,
4-(5-Carbonitrile-3-pyridylamino)-2-chloro-2'-methylbenzophenone oxime,
4-(3-Bromo-2-pyridylamino)-2-chloro-2'-methylbenzophenone oxime,
2-Chloro-2'-methyl-4-(8-quinolylamino)benzophenone oxime,
2-Chloro-4-(6-ethoxy-3-pyridylamino)-2'-methylbenzophenone oxime,
30 4-(4-Bromo-1-isoquinolylamino)-2-chloro-2'-methylbenzophenone oxime,
2-Chloro-2'-methyl-4-(2-methyl-5-trifluoromethyl-3-pyridylamino)benzophenone oxime,
2-Chloro-2'-methyl-4-(3-quinolylamino)benzophenone oxime,
2-Chloro-4-(4-ethoxy-3-pyridylamino)-2'-methylbenzophenone oxime,
2-Chloro-4-(2-ethoxy-3-pyridylamino)-2'-methylbenzophenone oxime,
35 2-Chloro-2'-methyl-4-(2-methyl-6-quinolylamino)benzophenone oxime,
2-Chloro-2'-methyl-4-(2-quinolylamino)benzophenone oxime,
2-Chloro-4-(7-chloro-4-quinolylamino)-2'-methylbenzophenone oxime,
2-Chloro-2'-methyl-4-(4-quinolylamino)benzophenone oxime,

- 2-Chloro-2'-methyl-4-(1-methyl-7-indolylamino)benzophenone oxime, and
 2-Chloro-2'-methyl-4-(1-methyl-5-indolylamino)benzophenone oxime,
 2-Chloro-2',5'-dimethyl-4-(4-methyl-3-pyridylamino)benzophenone oxime,
 2-Chloro-2',5'-dimethyl-4-(4-isoquinolylamino)benzophenone oxime,
 5 2-Chloro-4-(4-isoquinolylamino)-2',4',5'-trimethylbenzophenone oxime,
 2,3'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone oxime,
 2-Fluoro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone oxime,
 2,4'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone oxime,
 2-Chloro-4'-fluoro-4-(4-isoquinolylamino)-2'-methylbenzophenone oxime,
 10 4'-*n*-Butyl-2-chloro-4-(4-isoquinolylamino)-2'-methylbenzophenone oxime,
 2-Chloro-4-(5-isoquinolylamino)-2'-methylbenzophenone oxime,
 2-Chloro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone oxime,
 2-Fluoro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone oxime,
 2,4'-Dichloro-2'-methyl-4-(1-methyl-7-indolylamino)benzophenone oxime,
 15 2-Chloro-4-(1-methyl-7-indolylamino)-2',4',5'-trimethylbenzophenone oxime,
 2-Chloro-2',5'-dimethyl-4-(1-methyl-7-indolylamino)benzophenone oxime,
 2-Chloro-4-(3-ethoxy-4-isoquinolylamino)-2'-methylbenzophenone oxime,
 2-Chloro-4-(1-ethoxy-4-isoquinolylamino)-2'-methylbenzophenone oxime,
 4-(2-Benzoxazolylamino)-2-chloro-2'-methylbenzophenone oxime,
 20 4-(1-Methyl-2-benzimidazolylamino)-2-chloro-2'-methylbenzophenone oxime,
 4-(2-Benzothiazolylamino)-2-chloro-2'-methylbenzophenone oxime,
 2-Chloro-2'-methyl-4-(2-pyrimidylamino)benzophenone oxime,
 2-Chloro-2'-methyl-4-(7-methyl-purin-6-ylamino)benzophenone oxime,
 2-Chloro-2'-methyl-4-(2-methyl-5-benzothiazolylamino)benzophenone oxime,
 25 2-Chloro-2'-methyl-4-(pyrazin-2-ylamino)benzophenone oxime,
 2-Chloro-2'-methyl-4-(5-pyrimidylamino)benzophenone oxime,

and salts thereof with pharmaceutically acceptable acids, hydrates and solvates

- 30 Compounds of general formula Ic



35

wherein R₁, R₂, R₄, R₅, R₆, R₇, R₈ and X are as indicated above, and R₃ represents (C₁-C₃)alkyl, fluoro, chloro, bromo, methoxy, and hydroxy, and salts thereof with pharmaceutically acceptable acids, hydrates and solvates, are generally preferred.

- 5 In formula Ic, R₁ preferably represents methyl or halogen, more preferably F or Cl; R₂ represents one or more substituents, preferably hydrogen, halogen, (C₁-C₃)alkyl, methoxy or ethoxy, and R₃ represents methyl, methoxy or chloro.

Pharmacological methods

10

To study the effect of the compound of the present invention in vitro the inhibition of the IL-1 β and TNF- α secretion was measured using the following procedure:

15

Cytokine production was measured in the media from lipopolysaccharide (LPS) stimulated peripheral blood mononuclear cells. The mononuclear cells were isolated from human peripheral blood by Lymphoprep[®] (Nycomed, Norway) fractionation and suspended in RPMI 1640 (growth medium) with foetal calf serum (FCS, 2%), at a concentration of 5 x 10⁵ cells/ml. The cells were incubated in 24-well tissue culture plates in 1 mL aliquots. Test compounds were dissolved in dimethylsulfoxide (DMSO, 10 mM) and were diluted with the medium. Compounds were added to the cells for 30 minutes, then LPS (1 mg/mL final concentration) was added. The plates were incubated for 18 hours, and the concentration of IL-1 β and TNF- α in the medium was determined by enzyme-linked immunosorbent assays. The median inhibitory concentrations (IC₅₀) of the compounds were calculated. The results are shown in Table 1.

25

The compounds of the present invention also show similar activities in the ability to inhibit PMN (polymorphonuclear) superoxide secretion which is also indicative of potentially useful anti-inflammatory drugs. The compounds were tested using the following procedure:

30

Human polymorphonuclear (PMN) granulocytes were isolated from human blood by dextran sedimentation, Lymphoprep[®] fractionation and hypotonic lysis of contaminating erythrocytes.

35

Superoxide anion generation was measured as the superoxide dismutase inhibitable reduction of ferricytochrome C (Madhu, S.B. et al, Inflammation, 16, 241, (1992)). The cells were suspended in Hanks' balanced salt solution, and incubated for 10 minutes at

37°C with test compounds. The cells were primed by the addition of TNF- α (3 ng/mL final concentration) for 10 minutes, and then ferricytochrome C, (final concentration 750 μ g/mL), bovine serum albumin (BSA, final concentration 1 mg/mL) and formyl-methionyl-leucyl-phenylalanine (fMLP, final concentration 10^{-7} M) were added for 3 minutes. The cells were chilled on ice, and were spun down. The optical densities in the cell-free supernatant was measured in a spectrophotometer. The median inhibitory concentration (IC₅₀) of the compounds was calculated. The results, together with log P of the test compounds are shown in Table 1.

Table 1. Inhibition of cytokines and PMN-superoxide production in vitro by compounds of the present invention. The median inhibition concentration (IC₅₀, nM) Log P of compounds of the present invention

Comp. No.	IL-1 β	TNF- α	PMN-superoxide	
Comp. 114	31	5.0	15	4.2
Comp. 128	15.8	3.1	12.6	4.4
Ref. comp.	13	7.1	5.0	4.9

Ref. comp. : (4-(2-aminophenylamino)-2-chloro-2'-methylbenzophenone, Compound 156 disclosed in WO 98/32730.

These results show that the compounds of the present invention are able to inhibit the production of IL-1 β , TNF- α and PMN-superoxide, and showing a pharmacological activity comparable to a reference compound, thus making them potentially useful in the treatment of inflammatory diseases. The lower log P values of the compounds of the present invention reflects a lower lipophilicity, which indicate compounds with an improved bioavailability as compared to the reference compound.

p38 α MAP kinase assay

Cell culture

COS-1 cells (derived from African green monkey kidney fibroblast-like cell containing wild-type T antigen under control of the SV40 promotor) were obtained from ATCC (ATCC no. CRL-1650) and grown in growth medium (DMEM without phenolred, 10% FCS, 2 mM L-

glutamine, 100U penicillin and 100 µg streptomycin/ml) at 37°C with 5% CO₂. The cells were passaged twice a week by trypsinization (0.25% trypsin, 1 mM EDTA in PBS) and were split 1:10. The medium was changed every second or third day. The cell line was regularly tested with the Mycoplasma PCR Primer Set (Stratagene) and found to be free of

5 *Mycoplasma*. Tissue culture media, FCS, L-glutamine and penicillin and streptomycin are from Bribco BRL, Gaithersburg, MD, USA.

Transient expression of COS-1 cells

On day one COS-1 cells were seeded in 143 cm² petridish with a density of 2x10⁴cells/cm²
10 In growth medium. At day 2 the cells were co-transfected with 5 µg (total) of experimental plasmid DNA, expressing the FLAG-p38α and FLAG-MKK6(EE). The plasmids were introduced into the COS-1 cells in serum-free medium using DOTAP™ (Boehringer-Mannheim, Mannheim, Germany). Plasmid DNA was prepared and purified using the QIAGEN EndoToxin-free Maxiprep-500 kit (Hilden, Germany). Briefly, DNA and DOTAP™
15 were mixed for exactly 15 min. at 37°C in the CO₂ incubator. The transfection-mixture was hereafter transferred to a 15-mL falcon-tube and transfection-medium (DMEM with L-Glutamine and Pen./Strep. but without serum) was added to the transfection-mixture, followed by addition to the cell-monolayer. After 4 hours of incubation with DOTAP™ and plasmids, the medium containing double amount of serum was added to the cells bringing
20 the final concentration of serum up to 10%. The cells were then incubated for 24 hours before kinase reaction.

Immunoprecipitation

After 24 hrs of incubation the reaction was stopped by putting the petridish on an ice-
25 bath. The medium was aspirated, and the cell monolayer was washed once in ice-cold PBS (137 mM NaCl, 1.5 mM KH₂PO₄, 2.7 mM KCl, 8.1 mM Na₂HPO₄·2H₂O), and hereafter solubilised for 10 min. in 1.5 mL lysis buffer (50 mM HEPES, pH 7.5, 150 mM NaCl, 10 mM EDTA, 10 mM Na₄P₂O₇, 100 mM NaF, 2 mM Na₃VO₄, 1% Triton-X-100, Pefabloc 500 µM, Leupeptin 10 µg/µl, Aprotinin 10 µg/µl) was added. The cell-monolayer was scraped by a
30 rubber-policeman, and transferred to an Eppendorf tube. The solubilised cells were clarified by centrifugation at 10.000xg for 10 min. at 4°C. The supernatant was transferred to 50 µl prewashed Protein G Sepharose beads in HNT-buffer (30 mM HEPES, pH 7.5, 30 mM NaCl, 0.1% Triton X-100) and were incubated with 2 µg/sample of monoclonal anti-FLAG™ M2 antibody (raised against the FLAG-epitope, NH₂-Asp-Tyr-Lys-Asp-Asp-Asp-Asp-
35 Lys-COOH) for 1 hour at room temperature. The anti-FLAG M2 monoclonal antibody was obtained from Sigma (cat. no. F-3165). Approx. 60 µg protein of clarified cell lysate were added to the preadsorbed anti-FLAG™ antibodies on Protein G Sepharose beads and incubated for 90 min. at 4°C in a blood sample mixer. After the immunoprecipitation

period the Sepharose beads were washed twice in lysis buffer and twice in a kinase reaction buffer (25 mM HEPES pH 7.5, 10 mM magnesium acetate, 50 μ M ATP).

Incubation of the compounds with purified p38 α kinase

The pre-washed Immunoprecipitated anti-FLAG-p38 adsorbed on Protein G Sepharose beads was washed twice in 1 \times kinase-buffer (25 mM HEPES pH 7.5, 10 mM magnesium acetate, 50 μ M ATP), and the supernatant was aspirated. The compounds were diluted in 1 \times kinase buffer at the appropriate concentration. The compounds were added to the washed Immunoprecipitated and activated FLAG-p38 adsorbed on the Protein G Sepharose beads for 30 min. at 30°C in a volume of 100 μ l. Every 10 min. the Eppendorf tubes were tapped to ensure that the beads and the compounds were in the solution. After 30 min. incubation, the beads were spinned down and the supernatant was aspirated.

p38 α MAP Kinase Reaction

The kinase reaction was started by adding 1 μ g GST-ATF-2 substrate (Santa Cruz, LaJolla, CA, USA, cat. no. sc-4114) together with 2 μ Ci γ -³²P-ATP in 1 \times kinase-buffer per sample. The reaction was allowed to proceed for 30 min. at 30°C, and it was stopped by adding 40 μ l of 2 \times SDS-sample buffer to the kinase reaction. The samples were boiled, spinned down, and resolved on a 15% SDS-PAGE. The dried SDS-PAGE gel was exposed to a Phospho-Imager screen and the radioactive PHAS-1 bands were quantified by the STORM860 Phospho-Imager (Molecular Dynamics, Sunnyvale, CA, USA) using the ImageQuANT software.

The inhibition exhibited by compound 114 in this assay is shown in table 2 below:

25

Table 2

Compound No.	IC ₅₀ values for the inhibition of p38 α MAP kinase, mean concentration (nM)
Compound 114	12.8
Reference compound, SB 203580	1056.6

Reference compound SB 203580 is a widely used reference compound for p38 α MAP kinase inhibition. The compound is commercially available at Calbiochem (Calbiochem-Novabiochem LaJolla, CA, USA)

30

These results show, that the compounds of this invention are potent p38 α MAP kinase inhibitors with an improved pharmacological activity compared to a reference compound, thus making them potentially useful in the treatment of inflammatory diseases.

5

To study the compounds of the present invention *in vivo* the 12-*O*-tetradecanoylphorbol-13-acetate (TPA) induced murine chronic skin inflammation model can be used (De Young, L.M. et al., *Agents Actions* 26, 335-341 (1989); Carlson, R.P. et al., *Agents Actions* 17, 197-204 (1985); Alford, J.G. et al., *Agents Action* 37, (1992); Stanley, P.L. et al., *Skin Pharmacol.* 4, 262-271 (1991)), cf. the description of method in WO 98/32730 hereby incorporated by reference. These results show that compounds of the present invention are of the same potency compared to known reference compounds, e.g. hydrocortisone with its known side effects, whereas the compounds of the present invention are well tolerated and are non-toxic. Some members of the present class of compounds show a very low absorption, thus making them especially useful in the treatment of various dermatological diseases. In general, they may be administered by e.g. oral, intravenous, intranasal, topically or transdermal routes.

10
15

Methods of preparing compounds of formula I

20

The compounds of the present invention may be prepared in a number of ways well known to those skilled in the art of organic synthesis. The compounds of the present invention may be synthesised using the methods outlined below, together with methods known in the art of synthetic organic chemistry, or variations thereof as appreciated by those skilled in the art. Preferred methods include, but are not limited to, those described below.

25

The novel compounds of formula I, Ia, and Ib may be prepared using the reactions and techniques described in this section. The reactions are performed in solvents appropriate to the reagents and materials employed and are suitable for the transformations being effected. Also, in the synthetic methods described below, it is understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of experiment and work-up procedures, are chosen as standard conditions for that reaction, which should be readily recognised by one skilled in the art. It is understood by one skilled in the art of organic synthesis that the functionality present on various portions of the educt molecule must be compatible with the reagents and reactions proposed. Not all compounds of formula I falling into a given class may be compatible with some of the reaction conditions required in some of the methods described. Such restrictions to the substituents which are compatible with the reaction

30


35

conditions will be readily apparent to one skilled in the art and alternate methods can be used.

The following abbreviations have been used throughout this specification: BINAP= racemic or non-racemic 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, CDCl_3 =deuteriochloroform, DMF=N,N-dimethyl-formamide, DMSO-d_6 =hexadeuterodimethylsulfoxide, DMSO=dimethylsulfoxide, EtOAc = ethyl acetate, Et_2O = diethylether, $\text{Pd}_2(\text{dba})_3$ = tris(dibenzylideneacetone)dipalladium(0), *m*CPBA = *meta*-chloroperbenzoic acid, MeOH = methanol, NaOt-Bu = sodium-*tert*-butoxide, KOt-Bu =potassium-*tert*-butoxide

THF=tetrahydrofuran, TLC=thin layer chromatography.

SCHEME 1



L: F, Cl, Br, I, or OSO_2CF_3

Y: Cl, Br, I, $\text{OSO}_2\text{R}'$

FGI: Functional group interconversion

Compounds according to the present invention may be prepared by a process comprising coupling of an amine of the formula III with an bromide, iodide, fluoride, chloride or triflate with the formula II, as shown in Scheme 1, where R_1 , R_2 , R_3 , R_4 , R_5 , and X are as defined in general formula I, except that any substituents or functional group which are potentially reactive in the coupling reaction may themselves be protected before the coupling reaction is performed and subsequently removed.

The coupling reaction is carried-out using any of the methods for the formation of diphenylamines known to one skilled in the art of organic synthesis.

The preferred method is the palladium catalysed amination method which comprises coupling of an amine with an heteroarylhalogenide (or heteroaryltriflate) in the presence of a base, a suitable Pd source, and a suitable phosphine ligand in an inert solvent.

5 (References: Wolfe, J.P.; Wagaw, S.; Buchwald, S.L.; *J. Am. Chem. Soc.*, (1996), 118, 7215-16; Wagaw, S.; Buchwald, S. L.; *J. Org. Chem.*, (1996), 61, 7240-41; Wolfe, J.P.; Buchwald, S.L.; *Tetrahedron Lett.*, (1997), 38, 6359-62; Hong. Y. *et al.*, *Tetrahedron Lett.*, (1997), 38, 5607-10).

10 The palladium compound used in the process is not particularly limited, and as specific examples are palladium(II) acetate, palladium(II) chloride, palladium(II) bromide, dichlorobis(triphenylphosphine)palladium(II), tetrakis(triphenylphosphine)palladium(0), tris(dibenzylideneacetone)dipalladium(0). The preferred ligand include, but are not limited to, racemic or non-racemic 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (hereinafter
15 referred to as BINAP), tri-*o*-tolylphosphine, tri-*tert*-butylphosphine, 1,1'-bis(diphenylphosphino)-ferrocene, bis[(2-diphenylphosphino)phenyl]ether (DPEphos), 2-dicyclohexylphosphanyl-2'-dimethylaminobiphenyl, 2-(di-*tert*-butylphosphino)biphenyl, and 9,9-dimethyl-4,6-bis(diphenylphosphino)xanthene (Xantphos). The amount of palladium and ligand used in this process is typically in the range 0.1 to 10 % by mole
20 relative to the amount of the aromatic halide (or triflate) used.

Especially NaOt-Bu and caesium carbonate (Cs_2CO_3) have proven to be the best bases in this process, but other bases may be used as well.

25 The reaction is typically performed at elevated temperature (80-120 °C) in inert solvents like 1,4-dioxane, toluene, benzene and tetrahydrofuran under an inert atmosphere like argon or nitrogen.

The coupling reaction may also be carried out by nucleophilic substitution of a
30 heteroarylhalogenide with an amine, either in the presence of a base in a polar aprotic solvent, or without a solvent. The preferred base is KOt-Bu or NaH, but other bases may be used as well. The preferred solvent is dimethyl sulfoxide, but other solvents such as DMF may be used as well. The reaction is carried out at elevated temperature (120°C - 150°C) for 12 - 24h.

35

Compounds according to the present invention in which R_4 is not hydrogen may be prepared by a process comprising coupling of an amine of the formula I ($\text{R}_4 = \text{H}$) with an

alkylating agent, as shown in Scheme 1, where R_1 , R_2 , R_3 , R_5 , and X are as defined in general formula I, except that any substituents or functional group which are potentially reactive in the coupling reaction may themselves be protected before the coupling reaction is performed and subsequently removed.

5

Typically alkylating agents of the general formula R-Y include, but are not limited to, iodides (Y=I), bromides (Y = Br), chlorides (Y = Cl) and sulfonates (Y=OSO₂R', where R' represents methyl, trifluoromethyl or 4-methylphenyl).

10 Compounds according to the present invention may in special cases be prepared by a simple functional group Interconversion (FGI), meaning a standard process, known to those skilled in the art of organic synthesis, where a functional group in compounds with the general formula I is transformed into a different functional group in one or more synthetic steps, leading to a new compound with the general formula I. Examples of such
15 processes are, but are not limited to, hydrolysis of an ester to give an acid under basic conditions; deprotection of an methylether to give an phenol by treatment with e.g. borontribromide (BBr₃); catalytic hydrogenation of an olefin to give an saturated hydrocarbon and reduction of a nitro group to give an amine.


20 Compounds according to the present invention with the general formula I were X = S may be prepared from the ketone (with the general formula I, C =O) by such a FGI process, by using one of the many thiocarbonylating reagent, known to those skilled in the art of organic synthesis. Examples of such thiocarbonylating reagents include, but are not limited to, phosphorous pentasulfide (P₄S₁₀), or Lawesson's reagent (2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiaphosphetane-2,4-disulfide) or the like.

25 Compounds accordingly to the present invention with the general formula I were X = N-OH may be prepared from the ketone (with the general formula I, C =O) by treatment with hydroxylamine, or a protected derivative thereof followed by deprotection, in an appropriate solvent like e.g. pyridine or methanol.

30

35

SCHEME 2



hal: Br, I

- Compounds according to the present invention with the general formula III may be prepared by several methods known to those skilled in the art of organic synthesis. One useful sequence is shown in Scheme 2. The key step comprising coupling of a bromide (or iodide) with the general formula VI with an acid chloride with the general formula V to afford the benzophenone with the general formula IV. This compound IV may then be reduced to the corresponding amine with the general formula III by treatment with standard reducing agents. Examples of such reducing agents include, but are not limited to, stannous chloride dihydrate; hydrogen, ammonium formate, or hydrazine hydrate and a catalytic amount of palladium on carbon. The coupling reaction is done by transforming the bromide (VI) into a reactive organometallic intermediate, e.g. by treatment with butyllithium to afford the lithium derivative or by treatment with magnesium to afford the magnesium derivative. The reactivity of this intermediate is then modulated by transmetalation to e.g. zinc, by treatment with ZnCl_2 , ZnBr_2 , or ZnI_2 . This organozinc compound is then coupled with the acid chloride, with the general formula V, under the

influence of a palladium(0) complex in catalytic amount. Examples of such catalyst include but are not particularly limited to tetrakis(triphenylphosphine)palladium(0), tetrakis(triphenylarsine)palladium(0), dichlorobis(triphenylphosphine)palladium(II), or benzylchlorobis(triphenylphosphine)palladium(II).

5

As shown in Scheme 2 compounds with the general formula IV (X = O) may be transformed by a FGI process to give compounds with the general formula IV (X = S or X = N-OH) as described above. This is only to illustrate the flexibility in the synthesis and in general the described sequence of processes is only one of many possible strategies for the synthesis of compound of the present invention. That is, it may be more advantageous in some cases to alter the sequence of the processes described above. The described sequence of processes is not considered as being limited for the preparation of the compounds of the present invention with the general formula I and alteration of the reaction sequence is an obvious alternative for those skilled in the art of organic synthesis.

10

Pharmaceutical compositions

15

In another aspect, the invention relates to a pharmaceutical composition comprising, as an active component, a compound of formula I, Ia, Ib or Ic together with a pharmaceutically acceptable excipient or carrier. The term "pharmaceutically acceptable" is intended to indicate that the excipient or carrier included in the composition is compatible with the other ingredients and not toxic or otherwise deleterious to a patient to whom the composition is administered.

20

25

Pharmaceutical compositions of the invention may be in unit dosage form such as tablets, pills, capsules, powders, granules, elixirs, syrups, emulsions, ampoules, suppositories or parenteral solutions or suspensions; for oral, parenteral, ophthalmic, transdermal, intra-articular, topical, pulmonal, nasal, buccal or rectal administration or in any other manner appropriate for the formulation of anti-inflammatory compounds and in accordance with accepted practices such as those disclosed in *Remington: The Science and Practice of Pharmacy*, 19th Ed., Mack Publishing Company, 1995. The term "unit dosage" is intended to indicate a unitary, i.e. a single dose which is capable of being administered to a patient, and which may be readily handled and packed, remaining as a physically and chemically stable unit dose comprising either the active component as such or a mixture of it with solid or liquid pharmaceutical excipients or carriers. In the composition of the invention, the active component may be present in an amount of from about 0.1-100% by weight of the composition.

30

35

For oral administration in the form of a tablet or capsule, a compound of formula I may suitably be combined with an oral, non-toxic, pharmaceutically acceptable carrier such as ethanol, glycerol, water or the like. Furthermore, suitable binders, lubricants, disintegrating agents, flavouring agents and colourants may be added to the mixture, as appropriate. Suitable binders include, e.g., lactose, glucose, starch, gelatin, acacia gum, tragacanth gum, sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes or the like. Lubricants include, e.g., sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride or the like. Disintegrating agents include, e.g., starch, methyl cellulose, agar, bentonite, xanthan gum or the like.

For the preparation of solid compositions such as tablets, the active compound of formula I is mixed with one or more excipients, such as the ones described above, and other pharmaceutical diluents such as water to make a solid preformulation composition containing a homogenous mixture of a compound of formula I. The term "homogenous" is understood to mean that the compound of formula I is dispersed evenly throughout the composition so that the composition may readily be subdivided into equally effective unit dosage forms such as tablets or capsules. The preformulation composition may then be subdivided into unit dosage forms containing from about 0.05 to about 1000 mg, in particular from about 0.1 to about 500 mg, of the active compound of the invention.

Liquid formulations for either oral or parenteral administration of the compound of the invention include, e.g., aqueous solutions, syrups, aqueous or oil suspensions and emulsion with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil. Suitable dispersing or suspending agents for aqueous suspensions include synthetic or natural gums such as tragacanth, alginate, acacia, dextran, sodium carboxymethylcellulose, gelatin, methylcellulose or polyvinylpyrrolidone.

For parenteral administration, e.g. intramuscular, intraperitoneal, subcutaneous or intravenous injection or infusion, the pharmaceutical composition preferably comprises a compound of formula I dissolved or solubilised in an appropriate, pharmaceutically acceptable solvent. For parenteral administration, the composition of the invention may include a sterile aqueous or non-aqueous solvent, in particular water, isotonic saline, isotonic glucose solution, buffer solution or other solvent conventionally used for parenteral administration of therapeutically active substances, in particular antiproliferative agents. The composition may be sterilised by, for instance, filtration through a bacteria-retaining filter, addition of a sterilising agent to the composition, irradiation of the composition, or heating the composition. Alternatively, the compound of

the invention may be provided as a sterile, solid preparation, e.g. a freeze-dried powder, which is dissolved in sterile solvent immediately prior to use.

5 The composition intended for parenteral administration may additionally comprise conventional additives such as stabilisers, buffers or preservatives, e.g. antioxidants such as methylhydroxybenzoate or the like.

10 Compositions for rectal administration may be in the form of a suppository incorporating the active ingredient and a carrier such as cocoa butter, or in the form of an enema.

15 Compositions suitable for intra-articular administration may be in the form of a sterile aqueous preparation of the active ingredient which may be in microcrystalline form, for example, in the form of an aqueous microcrystalline suspension. Liposomal formulations or biodegradable polymer systems may also be used to present the active ingredient for both intra-articular and ophthalmic administration.

20 Compositions suitable for topical administration, including eye treatment, include liquid or semi-liquid preparations such as liniments, lotions, gels, applicants, oil-in-water or water-in-oil emulsions such as creams, ointments or pastes; or solutions or suspensions such as drops. For topical administration, the active component preferably comprises from 1% to 20% by weight of the composition, but the active ingredient may comprise as much as 50% w/w.

25 Compositions suitable for administration to the nasal or buccal cavity or for inhalation include powder, self-propelling and spray formulations, such as aerosols and atomizers. Compositions suitable for nasal or buccal administration may comprise 0.1% to 20% w/w, for example about 2% w/w of active ingredient.

30 The composition may additionally comprise one or more other active components conventionally used in the treatment of various inflammatory diseases and conditions. Examples of such additional active components may be selected from the group consisting of glucocorticoids, vitamin D and vitamin D analogues, antihistamines, platelet activating factor (PAF) antagonists, anticholinergic agents, methylxanthines, β -adrenergic agents, COX-2 inhibitors, salicylates, infomethacin, flufenamate, naproxen, timegadine, gold salts, 35 penicillamine, serum cholesterol lowering agents, retinoids, zinc salts and salicylazosulfapyridine.

In a further aspect, the invention relates to the use of a compound of formula I for the preparation of a medicament for the treatment or prophylaxis of inflammatory diseases or conditions. In a still further aspect, the invention relates to a method of treating inflammatory diseases or conditions, the method comprising administering, to a patient in
5 need thereof, an effective amount of a compound of formula I.

A suitable dosage of the compound of the invention will depend, inter alia, on the particular compound selected for the treatment, the route of administration, the age and condition of the patient, the severity of the disease to be treated and other factors well
10 known to the practising physician. The compound may be administered either orally or parenterally according to different dosing schedules, e.g. daily or with weekly intervals. In general a single dose will be in the range from 0.01 to 400 mg/kg body weight, such as from 0.1 to 100 mg/kg body weight. The compound may be administered as a bolus (i.e. the entire daily dose is administered at once) or in divided doses two or more times a day.

15 Inflammatory diseases or conditions contemplated for treatment with the present compounds are inflammatory diseases where modulation of cytokine expression and secretion may be mediated by MAP kinases such as the p38 MAP kinase as discussed above. Examples of inflammatory diseases or conditions believed to be mediated by the
20 p38 MAP kinase are selected from the group consisting of asthma, allergy, arthritis, including rheumatoid arthritis, osteoarthritis and spondyloarthritis, gout, atherosclerosis, inflammatory bowel disease, Crohn's disease, proliferative and inflammatory skin disorders, such as psoriasis, atopic dermatitis and acne vulgaris, uveitis, sepsis, septic shock, AIDS related diseases and osteoporosis.

25 The treatment may additionally involve administration of one or more other anti-inflammatory active components such as glucocorticoids, vitamin D and vitamin D analogues, antihistamines, platelet activating factor (PAF) antagonists, anticholinergic agents, methylxanthines, β -adrenergic agents, COX-2 inhibitors, salicylates, infomethacin,
30 flufenamate, naproxen, timegadine, gold salts, penicillamine, serum cholesterol lowering agents, retinoids, zinc salts and salicylazosulfapyridine.

The invention is further described in the following general procedures, preparations and examples which are not in any way intended to limit the scope of the invention as claimed.




35






EXAMPLES







General procedures, preparations and Examples




Specific examples of compounds of formula I are listed in Table 3. All melting points are uncorrected. For ^1H and ^{13}C nuclear magnetic resonance (NMR) spectra (300 MHz for ^1H) chemical shift values (δ) (in ppm) are quoted, unless otherwise specified, for deuteriochloroform and hexadeuterodimethyl-sulfoxide solutions relative to internal tetramethylsilane (δ 0.00) or chloroform (^1H NMR δ 7.25, ^{13}C NMR δ 76.81). The value for a multiplet (m), eventually defined as a doublet (d), triplet (t) or quartet (q) is given at the approximate mid point unless a range is quoted. A signal may also be defined as a singlet (s) or broad singlet (b). The organic solvents used were anhydrous. The term "chromatography" refers to column chromatography using the flash technique and was performed on silica gel.







Table 3 Compounds of General formula I





Comp. No. Example No.	X	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇
Comp. 101 Ex. 1	O	CH ₃	H	2-Cl	H		H	-
Comp. 102 Ex. 2	O	CH ₃	H	2-Cl	H		H	-
Comp. 103 Ex. 3	O	CH ₃	H	2-Cl	H	Same as ex.2	5-NO ₂	-
Comp. 104 Ex. 4	O	CH ₃	H	2-Cl	H	Same as ex.2	5-NO ₂ ,6-NH ₂	-
Comp. 105 Ex. 5	O	CH ₃	H	2-Cl	H	Same as ex.2	4-COOH,6-Cl	-
Comp. 106 Ex. 6	O	CH ₃	H	2-Cl	H	Same as ex.2	3-CN	-
Comp. 107 Ex. 7	O	CH ₃	H	2-Cl	H		H	-
Comp. 108 Ex. 8	O	CH ₃	H	2-Cl	H	Same as ex.2	5,6-di-NH ₂	-
Comp. 109 Ex. 9	O	CH ₃	H	2-Cl	H	Same as ex.2	3-NO ₂	-
Comp. 110	O	CH ₃	H	2-Cl	H	Same as ex.2	3-NH ₂	-

Comp. No. Example No.	X	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇
Ex. 10								
Comp. 111	O	CH ₃	H	2-Cl	H	Same as ex.2	5-NH ₂	-
Ex. 11								
Comp. 112	O	CH ₃	H	2-Cl	H		H	-
Ex. 12								
Comp. 113	O	CH ₃	H	2-Cl	H		H	-
Ex. 13								
Comp. 114	O	CH ₃	H	2-Cl	H		H	H
Ex. 14								
Comp. 115	O	CH ₃	H	2-Cl	H		3-COOC ₄ H ₉	-
Ex. 15								
Comp. 116	O	CH ₃	H	2-Cl	H	Same as ex.7	4-CH ₃	-
Ex. 16								
Comp. 117	O	CH ₃	H	2-Cl	H	Same as ex.7	6-CH ₃	-
Ex. 17								
Comp. 118	O	CH ₃	H	2-Cl	H	Same as ex.7	5-Br	-
Ex. 18								
Comp. 119	O	CH ₃	H	2-Cl	H	Same as ex.7	5-CN	-
Ex. 19								
Comp. 120	O	CH ₃	H	2-Cl	H	same as ex.2	3-Br	-
Ex. 20								
Comp. 121	O	CH ₃	H	2-Cl	H		H	H
Ex. 21								
Comp. 122	O	CH ₃	H	2-Cl	H	same as ex.7	6-OC ₂ H ₅	-
Ex. 22								

Comp. No. Example No.	X	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇
Comp. 123 Ex. 23	O	CH ₃	H	2-Cl	H		4-Br	H
Comp. 124 Ex. 24	O	CH ₃	H	2-Cl	H	same as ex.7	2-CH ₃ , 5-CF ₃	-
Comp. 125 Ex. 25	O	CH ₃	H	2-Cl	H		H	H
Comp. 126 Ex. 26	O	CH ₃	H	2-Cl	H	same as ex.7	4-OC ₂ H ₅	-
Comp. 127 Ex. 27	O	CH ₃	H	2-Cl	H	same as ex.7	2-OC ₂ H ₅	-
Comp. 128 Ex. 28	O	CH ₃	H	2-Cl	H		H	H
Comp. 129 Ex. 29	O	CH ₃	H	2-Cl	H		H	2-CH ₃
Comp. 130 Ex. 30	O	CH ₃	H	2-Cl	H		H	7-Cl
Comp. 131 Ex. 31	O	CH ₃	H	2-Cl	H		H	H
Comp. 132 Ex. 32	O	CH ₃	H	2-Cl	H	same as ex.30	H	H


Comp. No. Example No.	X	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇
Comp. 133 Ex. 33	O	CH ₃	H	2-Cl	H		H	H
Comp. 134 Ex. 34	O	CH ₃	H	2-Cl	H		H	H
Comp. 135 Ex. 35	O	CH ₃	5-CH ₃	2-Cl	H	Same as ex. 7	4-CH ₃	-
Comp. 136 Ex. 36	O	CH ₃	5-CH ₃	2-Cl	H	Same as ex. 14	H	H
Comp. 137 Ex. 37	O	CH ₃	4-CH ₃ , 5-CH ₃	2-Cl	H	Same as ex. 14	H	H
Comp. 138 Ex. 38	O	CH ₃	3-Cl	2-Cl	H	Same as ex. 7	4-CH ₃	-
Comp. 139 Ex. 39	O	CH ₃	H	2-F	H	Same as ex. 7	4-CH ₃	-
Comp. 140 Ex. 40	O	CH ₃	4-Cl	2-Cl	H	Same as ex. 7	4-CH ₃	-
Comp. 141 Ex. 41	O	CH ₃	4-F	2-Cl	H	Same as ex. 14	H	H
Comp. 142 Ex. 42	O	CH ₃	4-n-butyl	2-Cl	H	Same as ex. 14	H	H
Comp. 143 Ex. 43	O	CH ₃	H	2-Cl	H		H	H
Comp. 144 Ex. 44	O	CH ₃	4-OC H ₃	2-Cl	H	Same as ex. 14	H	H
Comp. 145 Ex. 45	O	CH ₃	4-OC	2-F	H	Same as ex. 14	H	H

Comp. No. Example No.	X	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇
			H ₃					
Comp.146 Ex. 46	O	CH ₃	4-Cl	2-Cl	H	Same as ex. 33	H	H
Comp.147 Ex. 47	O	CH ₃	4- CH ₃ ,	2-Cl	H	Same as ex. 33	H	H
Comp.148 Ex. 48	O	CH ₃	5- CH ₃	2-Cl	H	Same as ex. 33	H	H
Comp.149 Ex. 49	O	CH ₃	H	2-Cl	H	Same as ex. 14	3-OC ₂ H ₅	H
Comp.150 Ex. 50	O	CH ₃	H	2-Cl	H	Same as ex. 14	1-OC ₂ H ₅	H
Comp.151 Ex. 51	O	CH ₃	H	2-Cl	H		-	-
Comp.152 Ex. 52	O	CH ₃	H	2-Cl	H		-	-
Comp.153 Ex. 53	O	CH ₃	H	2-Cl	H		-	-
Comp.154 Ex. 54	O	CH ₃	H	2-Cl	H		-	-
Comp.155 Ex. 55	O	CH ₃	H	2-Cl	H		-	-
Comp.156 Ex. 56	O	CH ₃	H	2-Cl	H		-	-

Comp. No, Example No.	X	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇
Comp.157 Ex. 57	O	CH ₃	H	2-Cl	H		-	-
Comp.158 Ex. 58	O	CH ₃	H	2-Cl	H		-	-
Comp. 159 Ex. 59	O	CH ₃	H	2-Cl	H		-	-
Comp. 160 ex. 60	O	CH ₃	H	2-Cl	H		-	-

The numbering in Table 2 refers to the numbering in the formula below, R₆ and R₇ being defined as for general formula I:

5



10

General procedure 1

Coupling of compounds of the general formula II with compounds of the general formula
15 III to give compounds of the general formula I, or a protected derivative thereof.

In a screw cap vessel an amine (1.0 eq.) with the general formula III was dissolved in 1,4-dioxane or toluene, and a halogenide (1.0 - 1.1 eq.), with the general formula II was added. The vessel was flushed with argon, the base (Cs₂CO₃ or NaOt-Bu, 1.4 eq.),
20 Pd₂(dba)₃ (0.02 eq.), and BINAP (0.04 eq.) were added, the vessel was again flushed with argon and closed. The resulting suspension was first shaken vigorously at room

temperature for 5 min and then at 100 °C - 110°C for 4-20 h or until the halogenide II had disappeared as seen on TLC. The reaction mixture was allowed to cool to room temperature. The reaction mixture was poured into a mixture of EtOAc and water. The organic phase was separated, and the water phase was extracted with EtOAc three times.
5 The combined organic phases were washed with water and saturated aqueous NaCl, dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified either by chromatography and/or crystallisation to afford the coupled product with the general formula I, or a protected derivative thereof.

10 General procedure 2

Coupling of compounds of the general formula II with compounds of the general formula III to give compounds of the general formula I, or a protected derivative thereof.

15 An amine of the general formula (III) (1 eq.) was dissolved in DMSO (0.1 - 0.2M), and a heteroarylchloride or heteroarylfluoride (1 eq.) of the general formula (II) was added. Potassium *t*-butoxide, 2 eq., or 3 eq. if compound II was provided as the hydrochloride, was dissolved in DMSO (0.2 - 0.4M), and the solution was added dropwise to the reaction mixture. The reaction was stirred under an argon atmosphere at 120 - 150°C for 12 - 24 h. The reaction mixture was poured into water and extracted with EtOAc three times. The
20 combined organic phases were washed with saturated aq. NaCl, dried (MgSO₄), filtered and concentrated *in vacuo*, and the crude product was purified by flash chromatography to afford the coupled product with the general formula I, or a protected derivative thereof.

General procedure (3)

25 Coupling of compounds of the general formula VI with compounds of the general formula V to give compounds of the general formula IV, or a protected derivative thereof.

The bromide (80 mmol) with the general formula VI was dissolved in dry THF (65 mL) and cooled with stirring to -78 °C under an atmosphere of argon. *n*-Butyllithium (80 mmol, 1.6
30 M solution in hexane) was then added dropwise, keeping the internal temperature below -65 °C and stirring the resulting mixture for further 15 min. A THF solution of ZnCl₂ (100 mmol, 1.0 M) was added dropwise and the reaction mixture was allowed to warm to room temperature. After 2 h the reaction mixture was cooled to 0 °C, and tetrakis(triphenylphosphine)palladium(0) (4.0 mmol) was added followed by the dropwise
35 addition of the acid chloride (84 mmol), with the general formula V, in THF. The reaction mixture was allowed to warm to 20 °C and stirred for ca 16 h. The resulting yellow

solution, was filtered and the filtrate poured into a mixture of EtOAc/water 1:1, shaken and separated. The aqueous phase was extracted with two more portions of EtOAc. The organic phases were combined, dried (MgSO_4), filtered, and concentrated *in vacuo* to afford the crude product. Further purification was done by flash chromatography and/or crystallisation to give the title compound IV, or a protected derivative thereof.

General procedure (4)

Reduction of compounds of the general formula IV with stannous chloride dihydrate to give compounds of the general formula III, or a protected derivative thereof.

A mixture of a compound with the general formula IV (5 mmol) and stannous chloride dihydrate (5.64 g, 25 mmol) in absolute ethanol (50 mL) was heated to 70 °C under argon. After 1 hour, or until the starting material had disappeared as seen on TLC, the solution was allowed to cool to room temperature and then poured into ice/water. The pH was made alkaline by the addition of saturated sodium hydroxide (50 mL) before being extracted with ethyl acetate (3x100 mL). The organic phase was dried (MgSO_4), filtered and evaporated to afford the crude product. The crude product was further purified either by crystallisation or flash chromatography to yield the title compound or a protected derivative thereof.

General procedure (5)

Oxidation of compounds of the general formula I with *m*CPBA to give the corresponding N-oxides

A compound of the general formula I (1 eq.) was dissolved in CH_2Cl_2 and *m*CPBA (1.2 eq.) was added, and the reaction mixture was stirred at room temperature for 1 - 3h. $\text{Na}_2\text{S}_2\text{O}_5$ (1.5 eq.) was added, and the suspension was stirred and filtered. K_2CO_3 (3 eq.) was added to the filtrate and stirred for 0.5 - 1h. MgSO_4 was added, the suspension was filtered, and the filtrate was concentrated *in vacuo*. Purification by flash chromatography and/or by crystallisation.

Preparation 1:

2-Chloro-4-nitro-2'-methylbenzophenone, (Compound 1)

General procedure: 3

Starting compound VI: 2-Bromotoluene

Starting compound V: 2-Chloro-4-nitro-benzoyl chloride

Purification: Chromatography using EtOAc/pentane 1:9 as eluant

^{13}C NMR (CDCl_3): δ 195.1, 148.9, 145.5, 140.6, 135.0, 133.1, 132.7, 132.4, 131.9, 130.0, 125.9, 125.4, 121.9, 21.5

5 Preparation 2:

4-Amino-2-chloro-2'-methylbenzophenone, (Compound 2)

General procedure: 4

Starting compound IV: 2-Chloro-4-nitro-2'-methylbenzophenone (Compound 1)

Purification: Chromatography using EtOAc/pentane 1:9 followed by 1:4 as eluant

10 ^{13}C NMR (CDCl_3): δ 196.7, 150.5, 139.5, 137.6, 135.1, 133.9, 131.2, 130.7, 129.5, 127.5, 125.3, 116.0, 112.2, 20.3

Preparation 3:

2-Chloro-4'-methoxy-2'-methyl-4-nitrobenzophenone (compound 3)

15 General procedure: 3

Starting compound VI: 2-Bromo-5-methoxytoluene

Starting compound V: 2-Chloro-4-nitro-benzoyl chloride

Purification: Crystallization from a MeOH/cyclohexane 10:1

20 ^{13}C NMR (CDCl_3): δ 193.3, 163.3, 148.6, 146.3, 144.3, 135.5, 132.4, 129.6, 127.5, 125.3, 121.9, 118.1, 110.9, 55.5, 22.4

Preparation 4:

4-Amino-2-chloro-4'-methoxy-2'-methylbenzophenone (compound 4)

General procedure: 4

25 Starting compound IV: 2-Chloro-4'-methoxy-2'-methyl-4-nitrobenzophenone (compound 3)

Purification: Filtered through a short column of silica gel

^{13}C NMR (CDCl_3): δ 195.7, 161.7, 149.6, 141.6, 134.2, 133.3, 132.8, 131.4, 129.1, 117.0, 115.8, 112.4, 110.3, 55.3, 21.3

30

Preparation 5:

2,4'-Dichloro-2'-methyl-4-nitrobenzophenone (compound 5)

General procedure: 3

Starting compound VI: 2-Bromo-5-chlorotoluene

35 Starting compound V: 2-Chloro-4-nitro-benzoyl chloride

Purification: Chromatography using EtOAc/pentane 1:15 followed by 1:10 as eluant
 ^{13}C NMR (CDCl_3): δ 194.1, 149.0, 145.0, 142.6, 139.4, 133.5, 133.1, 132.7, 132.5, 130.1, 126.1, 125.5, 122.0, 21.4

5 Preparation 6:

4-Amino-2,4'-dichloro-2'-methylbenzophenone (compound 6)

General procedure: 4

Starting compound IV: 2,4'-Dichloro-2'-methyl-4-nitrobenzophenone (compound 5)

^{13}C NMR (CDCl_3): δ 195.4, 150.5, 139.8, 137.9, 136.5, 135.1, 133.7, 131.2, 130.9,
10 127.5, 125.6, 115.9, 112.3, 20.2

Preparation 7:

2-Chloro-4'-fluoro-2'-methyl-4-nitrobenzophenone (compound 7)

General procedure: 3

15 Starting compound VI: 2-Bromo-5-fluorotoluene

Starting compound V: 2-Chloro-4-nitro-benzoyl chloride

Purification: Chromatography using EtOAc/pentane 1:20 as eluant

^{13}C NMR (CDCl_3): δ 193.7, 165.1, 148.9, 145.3, 144.6, 134.7, 132.6, 131.4, 129.9,
20 125.5, 122.0, 119.5, 113.0, 21.8

Preparation 8:

4-Amino-2-chloro-4'-fluoro-2'-methylbenzophenone (compound 8)

General procedure: 4

Starting compound IV: 2-Chloro-4'-fluoro-2'-methyl-4-nitrobenzophenone (compound 7)

25 Purification: Filtered through a short column of silica gel

^{13}C NMR (CDCl_3): δ 195.4, 163.9, 150.3, 141.5, 135.5, 134.9, 133.5, 132.2, 128.0,
118.1, 115.9, 112.4, 112.3, 20.6

Preparation 9:

30 2-Fluoro-4-nitro-2'-methylbenzophenone (compound 9)

General procedure: 3

Starting compound VI: 2-Bromotoluene

Starting compound V: 2-Fluoro-4-nitro-benzoyl chloride

Purification: Crystallization from a mixture of MeOH/cyclohexane 6:1

35 ^{13}C NMR (CDCl_3): δ 193.1, 159.9, 150.3, 139.3, 136.4, 133.8, 132.6, 132.1, 131.7,
130.8, 125.8, 119.4, 112.4, 21.0

Preparation 10:

4-Amino-2-fluoro-2'-methylbenzophenone (compound 10)

General procedure: 4

- 5 Starting compound IV: 2-Fluoro-4-nitro-2'-methylbenzophenone (compound 9)

Purification: Filtered through a short column of silica gel

^{13}C NMR (CDCl_3): δ 194.3, 163.8, 153.0, 140.7, 136.1, 134.0, 130.8, 129.9, 127.9, 125.3, 116.8, 110.1, 101.3, 19.8

10 Preparation 11:

2-Chloro-2',5'-dimethyl-4-nitrobenzophenone (compound 11)

General procedure: 3

Starting compound VI: 2-Bromo-1,4-dimethylbenzene

Starting compound V: 2-Chloro-4-nitro-benzoyl chloride

- 15 Purification: Filtered through a short column of silica gel

^{13}C NMR (CDCl_3): δ 195.3, 148.9, 145.6, 137.4, 135.5, 135.0, 133.9, 132.8, 132.3, 132.2, 130.0, 125.5, 121.9, 21.1, 20.8

Preparation 12:

- 20 4-Amino-2-chloro-2',5'-dimethylbenzophenone (compound 12)

General procedure: 4

Starting compound IV: 2-Chloro-2',5'-dimethyl-4-nitrobenzophenone (compound 11)

Purification: Filtered through a short column of silica gel

- 25 ^{13}C NMR (CDCl_3): δ 196.8, 150.2, 139.3, 135.1, 134.9, 134.5, 133.8, 131.4, 131.1, 130.0, 128.0, 116.0, 112.2, 20.8, 19.9

Preparation 13:

2,3'-Dichloro-2'-methyl-4-nitrobenzophenone (compound 13)

General procedure: 3

- 30 Starting compound VI: 2-Bromo-3-chlorotoluene

Starting compound V: 2-Chloro-4-nitro-benzoyl chloride

Purification: Crystallization from a mixture of MeOH/cyclohexane 9:1

- ^{13}C NMR (CDCl_3): δ 194.6, 149.2, 144.5, 138.2, 137.3, 137.0, 133.5, 133.2, 130.7, 129.0, 126.5, 125.7, 122.0, 17.3

35

Preparation 14:

4-Amino-2,3'-dichloro-2'-methylbenzophenone (compound 14)

General procedure: 4

Starting compound IV: 2,3'-Dichloro-2'-methyl-4-nitrobenzophenone (compound 13)

Purification: Filtered through a short column of silica gel

^{13}C NMR (CDCl_3): δ 195.3, 150.9, 142.4, 135.8, 135.8, 134.8, 134.6, 131.0, 126.8,

5 126.7, 126.4, 116.2, 112.2, 17.1

Preparation 15:

2-Fluoro-4'-methoxy-2'-methyl-4-nitrobenzophenone (compound 15)

General procedure: 3

10 Starting compound VI: 2-Bromo-5-methoxytoluene

Starting compound V: 2-Fluoro-4-nitro-benzoyl chloride

Purification: Recrystallization from a mixture of MeOH/cyclohexane/ CH_2Cl_2

^{13}C NMR (CDCl_3): δ 191.3, 163.1, 159.3, 143.6, 135.0, 134.8, 131.2,

128.5, 119.4, 117.8, 112.2, 110.8, 55.5, 22.1

15

Preparation 16:

4-Amino-2-fluoro-4'-methoxy-2'-methylbenzophenone (compound 16)

General procedure: 4

Starting compound IV: 2-Fluoro-4'-methoxy-2'-methyl-4-nitrobenzophenone (compound

20 15)

Purification: Filtered through a short column of silica gel

^{13}C NMR (CDCl_3): δ 193.4, 163.1, 161.2, 152.2, 140.1, 133.7, 132.6, 131.7, 117.8,

116.6, 110.3, 110.1, 101.3, 55.3, 20.7

25 Preparation 17:

2-Chloro-4-nitro-2',4',5'-trimethylbenzophenone (compound 17)

General procedure: 3

Starting compound VI: 5-Bromo-1,2,4-trimethylbenzene

Starting compound V: 2-Chloro-4-nitro-benzoyl chloride

30 Purification: Recrystallisation from EtOAc

EO 01720-000 (CDCl_3)

^{13}C NMR (CDCl_3): δ 194.8, 148.7, 146.1, 143.0, 138.3, 134.1, 134.0, 133.4, 132.6,

132.5, 129.8, 125.4, 121.9, 21.2, 19.9, 19.2

35 Preparation 18:

4-Amino-2-chloro-2',4',5'-trimethylbenzophenone (compound 18)

General procedure: 4

Starting compound IV: 2-Chloro-4-nitro-2',4',5'-trimethylbenzophenone (compound 17)

Purification: Filtered through a short column of silica gel

^{13}C NMR (CDCl_3): δ 196.6, 149.9, 140.0, 136.6, 135.5, 134.7, 133.4, 132.7, 131.4, 128.5, 115.9, 112.2, 20.0, 19.7, 19.1

5

Preparation 19:

4'-*n*-Butyl-2-chloro-2'-methyl-4-nitrobenzophenone (compound 19)

General procedure: 3

Starting compound VI: 4-*n*-Butyl-2-methyliodobenzene

10

Starting compound V: 2-Chloro-4-nitro-benzoyl chloride

Purification: Chromatography, eluent EtOAc:Petroleumether 1:15

^{13}C NMR (CDCl_3): δ 194.6, 149.2, 148.7, 146.0, 141.1, 132.7, 132.7, 132.6, 132.3, 129.8, 125.9, 125.3, 121.8, 35.7, 33.1, 22.4, 21.8, 13.9

15

Preparation 20:

4-Amino-4'-*n*-butyl-2-chloro-2'-methylbenzophenone (compound 20)

General procedure: 4

Starting compound IV: 4'-*n*-Butyl-2-chloro-2'-methyl-4-nitrobenzophenone (compound 19)

20

Purification: Filtered through a short column of silica gel

^{13}C NMR (CDCl_3): δ 196.5, 150.0, 146.3, 138.3, 136.5, 134.7, 133.4, 131.5, 130.4, 128.4, 125.3, 115.9, 112.2, 35.6, 33.3, 22.4, 20.6, 13.9

Example 1: 2-Chloro-2'-methyl-4-(4-pyridylamino)benzophenone (Compound 101),

25

General procedure: 2

Starting compound II: 4-chloro-pyridine hydrochloride (0.30g, 2.0 mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.49g, 2.0 mmol)

Solvent: DMSO (15 mL)

30

Base: *t*-BuOK (0.67g, 6.0mmol) in DMSO (10 mL)

Reaction time: 22h

Reaction temperature: 150°C

Purification: Chromatography using EtOAc:MeOH 15:1

Product: Compound 101, 0.20g oil

35

^1H NMR (CDCl_3): δ 8.39(d,1H), 7.43(d,1H), 7.40-7.20(m,5H), 7.08(dd,1H), 6.97(d,3H), 2.50(s,3H)

Example 2: 2-Chloro-2'-methyl-4-(2-pyridylamino)benzophenone (Compound 102),

General procedure:2

Starting compound II: 2-chloro-pyridine (0.19 mL, 2.0mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.49g,
5 2.0mmol)

Solvent: DMSO (15mL)

Base: *t*-BuOK (0.45g, 4.0 mmol) in DMSO (10 mL)

Reaction time: 22 h

Reaction temperature: 150°C

10 Purification: Chromatography using EtOAc : petroleum ether 1:5

Product: Compound 102, 0.21g oil.

¹H NMR (CDCl₃): δ 8.28(d,1H), 7.65(d,1H), 7.58(t,1H), 7.41(d,1H), 7.40-7.25(m,4H),
7.19(t,1H), 6.95(s,1H), 6.86(m,2H), 2.47(s,3H)

15 **Example 3:** 2-Chloro-2'-methyl-4-(5-nitro-2-pyridylamino)benzophenone (Compound 103),

General procedure:2

Starting compound II: 2-chloro-5-nitro-pyridine (0.32g, 2.0 mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.49g,
20 2.0mmol)

Solvent: DMSO (10mL)

Base: *t*-BuOK (0.45g, 4.0mmol) in DMSO (20 mL)

Reaction time: 18h

Reaction temperature: 150°C

25 Purification: Chromatography using EtOAc : petroleum ether 2:7 - 1:3

Product: Compound 103, 0.09g crystalline compound

¹³C NMR (acetone d-6): δ 196.7, 159.4, 146.2, 144.2, 139.2, 139.0, 138.8, 133.8, 133.6,
133.5, 132.4, 132.4, 132.3, 131.0, 126.5, 121.0, 118.3, 112.1, 20.8

30 **Example 4:** 4-(6-Amino-5-nitro-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 104),

General procedure:2

Starting compound II: 2-amino-6-chloro-3-nitro-pyridine (0.69g, 4.0mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.98g,
35 4.0mmol)

Solvent: DMSO (15mL)

Base: *t*-BuOK (0.90g, 8.0mmol) in DMSO (25mL)

Reaction time: 48h

Reaction temperature: 150°C

Purification: Chromatography using CH₂Cl₂ : EtOAc 50:1 - 30:1

Product: Compound 104, 0.58g crystalline compound

¹³C NMR (DMSO): δ 195.9, 157.7, 154.7, 143.5, 138.0, 137.4, 135.7, 131.9, 131.8,

5 131.5, 131.4, 129.9, 125.8, 119.9, 119.5, 117.5, 103.2, 20.2

Example 5: 6-Chloro-2-(3-chloro-4-(2-methylbenzoyl)phenylamino) isonicotinic acid
(Compound 105),

General procedure:2

10 Starting compound II: 2,6-dichloro-4-cyano-pyridine (0.70g, 4.0mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.98g, 4.0mmol)

Solvent: DMSO (15 mL)

Base: *t*-BuOK (0.90g, 8.0mmol) in DMSO (25 mL)

15 Reaction time: 48h

Reaction temperature: 150°C

Purification: Chromatography using EtOAc:MeOH 5:1

Product: Compound 105, 0.52g oil, the cyano substituent has been hydrolysed during reaction or work up.

20 ¹³C NMR (DMSO): δ 195.7, 166.6, 154.8, 150.5, 147.2, 145.0, 138.4, 137.0, 132.3, 132.1, 131.2, 131.1, 129.5, 129.4, 125.7, 118.0, 115.6, 114.8, 111.3, 20.0

Example 6: 4-(6-carbonitrile-2-pyridylamino)-2-chloro-2'-methylbenzophenone
(Compound 106),

25 General procedure:2

Starting compound II: 2-chloro-3-cyano-pyridine (0.28g, 2.0 mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.49g, 2.0 mmol)

Solvent: DMSO (10 mL)

30 Base: *t*-BuOK (0.45g, 4.0mmol) in DMSO (20 mL)

Reaction time: 18h

Reaction temperature: 150°C

Purification: Chromatography using EtOAc : petroleum ether 1:1 - 1:0

Product: Compound 106, 0.07g crystalline compound

35 ¹³C NMR (CDCl₃): δ 195.9, 155.5, 144.3, 142.9, 142.1, 140.3, 140.0, 136.0, 133.2, 132.6, 132.1, 131.8, 131.1, 128.8, 125.7, 125.3, 115.7, 105.7, 102.4, 21.3

Example 7: 2-Chloro-2'-methyl-4-(3-pyridylamino)benzophenone (Compound 107),

General procedure: 1

Starting compound II: 3-chloro-pyridine (1.18 mL, 12.0mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (2.95g,

5 12.0mmol)

Solvent: Toluene (60 mL)

Base: *t*-BuONa

Reaction time: 45h

Reaction temperature: 110°C

10 Purification: Chromatography using CH₂ Cl₂ : acetone 4:1

Product: Compound 107, 2.05g crystalline compound

¹³C NMR (CDCl₃): δ 196.5, 146.8, 144.4, 142.7, 138.7, 138.1, 137.3, 134.9, 133.3,

131.4, 131.1, 130.4, 129.9, 127.1, 125.4, 124.0, 117.0, 113.4, 20.5

15 **Example 8:** 2-Chloro-4-(5,6-diamino-2-pyridylamino)-2'-methylbenzophenone (Compound 108),

Compound 104 (0.38g, 1.0mmol) was dissolved in EtOH (10mL) and DMF (10mL). SnCl₂

*H₂O (1.12g, 5.0mmol) was added and the solution was stirred at 90°C for 20h. The

reaction mixture was poured into ice water (100mL), 2 N NaOH (50mL) was added, and

20 the water phase was extracted with EtOAc (50mL) and Et₂O (2 x 50 mL). The combined organic phases were washed with saturated aqueous NaCl (50 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo*. The residue was purified by flash chromatography EtOAc : petroleum ether 2 : 1 - 1 : 0.

Product: Compound 108, 0.05g oil

25 ¹³C NMR (CDCl₃): δ 196.7, 148.9, 146.2, 145.6, 139.2, 137.9, 134.6, 133.1, 131.2, 130.8, 129.7, 129.2, 126.7, 125.4, 122.5, 117.4, 114.0, 102.1, 20.4

Example 9: 2-Chloro-2'-methyl-4-(3-nitro-2-pyridylamino)benzophenone (Compound 109),

30 2-Chloro-3-nitro-pyridine (0.32g, 2.0mmol) and 4-amino-2-chloro-2'-methylbenzophenone (compound 2) (0.49g, 2.0mmol) was mixed thoroughly. The reaction mixture was warmed at 100°C for 0.5h and then at 160°C for 0.5h. The reaction mixture was cooled to room temperature, and dissolved in saturated aqueous NaHCO₃ (10 mL) and EtOAc (30 mL). The phases were separated, and the organic phase was washed with

35 saturated aqueous NaHCO₃ (10mL), dried (MgSO₄), filtered and concentrated *in vacuo*.

Purification by flash chromatography using EtOAc: petroleum ether 1:7.

Product: Compound 109, 0.26g crystalline compound which was recrystallised from Et₂O

^{13}C NMR (CDCl_3): δ 196.6, 154.8, 149.2, 141.4, 138.9, 137.9, 135.6, 134.2, 133.4, 131.6, 131.6, 131.5, 130.6, 129.4, 125.5, 122.5, 119.0, 115.3, 20.9

Example 10: 4-(3-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 110),

Compound 109 (0.15g, 0.4mmol) was suspended in EtOH (10mL) and $\text{SnCl}_2 \cdot \text{H}_2\text{O}$ (0.46g, 2.0mmol) was added. The reaction mixture was stirred at 70°C for 1h, after which it was poured into ice water (50mL). 2N NaOH (50mL) was added and the water phase was extracted with EtOAc (3 x 20mL). The combined organic phases were dried (MgSO_4), filtered and concentrated *in vacuo*. Purification by flash chromatography using EtOAc:petroleum ether 2:3.

Product: Compound 110, 0.12g oil.

^{13}C NMR (CDCl_3): δ 196.9, 145.3, 143.9, 139.2, 138.9, 138.1, 134.2, 132.7, 131.7, 131.3, 131.0, 130.3, 130.0, 125.4, 124.6, 118.8, 118.5, 114.9, 20.5

Example 11: 4-(5-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 111),

Compound 103 (0.07g, 0.19mmol) was suspended in EtOH (5mL) and $\text{SnCl}_2 \cdot \text{H}_2\text{O}$ (0.21g, 1.0mmol) was added. The reaction mixture was stirred at 70°C for 3h and the reaction mixture was worked up as described for Compound 110. Purification by flash chromatography using EtOAc:petroleum ether 3:2.

Product: Compound 111, 0.038g oil

^{13}C NMR (CDCl_3): δ 196.7, 146.3, 146.0, 139.1, 137.9, 137.6, 135.1, 134.6, 133.1, 131.3, 130.9, 129.8, 129.6, 125.5, 125.4, 117.3, 113.9, 112.7, 20.5

Example 12: 2-Chloro-2'-methyl-4((2-pyridyl-N-oxide)amino)benzophenone (Compound 112)

General procedure: 5

Starting compound I: Compound 102 (0.16g, 0.5mmol)

Reaction time: 1h

Purification by chromatography using EtOAc:MeOH 8:1

Product: Compound 112, 0.098g crystalline compound which could be recrystallised from MeOH

^{13}C NMR (DMSO): δ 195.8, 146.0, 142.8, 137.7, 137.5, 132.2, 132.0, 131.8, 131.6, 131.4, 130.0, 127.0, 125.8, 121.0, 117.7, 116.1, 109.8, 20.2

Example 13: 2-Chloro-2'-methyl-4-((3-pyridyl-N-oxide)amino)benzophenone (Compound 113)

General procedure: 5

Starting compound I: Compound 107 (0.18g, 0.54 mmol)

5 Reaction time: 2h

Purification by chromatography using EtOAc:MeOH 7:1

Product: Compound 113, 0.12g crystalline compound

^{13}C NMR (CD_3OD): δ 198.5, 146.6, 143.7, 139.7, 139.4, 135.4, 134.0, 133.2, 133.0, 132.7, 132.6, 131.2, 130.4, 128.1, 126.8, 120.0, 119.6, 116.5, 20.8

10

Example 14: 2-Chloro-4-(4-isoquinolylamino)- 2'-methylbenzophenone (Compound 114),

General procedure: 1

Starting compound II: 4-bromo-isoquinoline (0.42g, 2.0mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.49, 2.0mmol)

15

Solvent: 1,4-dioxane (6mL)

Base: *t*-BuONa

Reaction time: 18h

Reaction temperature: 100°C

20

Purification: Chromatography using CH_2Cl_2 : acetone 10:1

Product: Compound 114, 0.64g amorphous compound.

^{13}C NMR (CDCl_3): δ 196.6, 150.1, 149.3, 139.0, 138.6, 137.9, 135.1, 133.5, 132.1, 131.3, 131.1, 130.9, 130.8, 129.7, 129.3, 129.3, 128.2, 127.9, 125.4, 121.6, 116.3, 112.6, 20.5

25

Example 15: *t*-Butyl 5-(3-chloro-4-(2-methylbenzoyl)phenylamino) nicotinoate (Compound 115,)

5-Bromonicotinic acid (4.7g, 23mmol) was added to 7 mL condensed isobutylene in CH_2Cl_2 (15mL), and afterwards conc. H_2SO_4 (0.75mL) was added. The reaction mixture was stirred in a screw cap vessel for 24h at room temperature, after which it was poured into saturated aqueous NaHCO_3 (100mL). The aqueous phase was extracted with EtOAc (100mL), the organic phase was washed with NaHCO_3 , dried (MgSO_4), filtered and concentrated. The crude product was purified by chromatography using the eluent EtOAc:petroleum ether 1:10.

30

35

Product: *t*-butyl 5-bromonicotinoate, crystalline compound.

^{13}C NMR (CDCl_3): δ 163.06, 154.02, 148.85, 139.38, 129.10, 120.48, 82.82, 28.12

General procedure: 1

Starting compound II: *t*-Butyl 5-bromonicotinoate, (0.12g, 0.45mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.13g, 0.53mmol)

Solvent: toluene (3mL)

5 Base: *t*-BuONa

Reaction time: 12h

Reaction temperature: 100°C

Purification: Chromatography using EtOAc:petroleum ether 1:2

Product: Compound 115, 0.035g oil

10 ¹³C NMR (CDCl₃): δ 196.7, 164.1, 146.1, 145.1, 144.8, 138.5, 138.2, 137.4, 134.8, 133.2, 131.5, 131.3, 131.0, 130.1, 128.4, 126.9, 125.5, 117.7, 113.7, 82.5, 28.1, 20.6

Example 16: 2-Chloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 116),

15 General procedure: 1

Starting compound II: 3-bromo-4-methyl pyridine (0.172g, 1.0mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2)(0.25g, 1.0mmol)

Solvent: 1,4-dioxane (5mL)

20 Base: *t*-BuONa

Reaction time: 18h

Reaction temperature: 110°C

Purification: Chromatography using EtOAc:petroleum ether 1:3 - 1:1

Product: Compound 116, 0.08g oil

25 ¹³C NMR (CDCl₃): δ 196.5, 148.5, 146.5, 146.1, 142.1, 139.0, 138.0, 135.4, 135.1, 133.5, 131.3, 131.0, 129.8, 129.3, 125.9, 125.4, 116.0, 112.3, 20.5, 17.5

Example 17: 2-Chloro-2'-methyl-4-(6-methyl-3-pyridylamino)benzophenone (Compound 117),

30 General procedure: 1

Starting compound II: 5-bromo-2-methyl pyridine (0.172g, 1.0mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g, 1.0 mmol)

Solvent: 1,4-dioxane (5mL)

35 Base: *t*-BuONa

Reaction time: 18h

Reaction temperature: 110°C

Purification: Chromatography using EtOAc:petroleum ether 1:4 - 1:2

Product: Compound 117, 0.15g oil

^1H NMR (CDCl_3): δ 8.37(d,1H), 7.45(dd,1H), 7.4-7.15(m,6H), 6.93(d,1H), 6.78(dd,1H), 6.29(s,1H), 2.54(s,3H), 2.44(s,3H)

5 Example 18: 4-(5-Bromo-3-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 118),

General procedure: 1

Starting compound II: 3,5-dibromo-pyridine (0.237g, 1.0 mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g,
10 1.0 mmol)

Solvent: 1,4-dioxane (5mL)

Base: *t*-BuONa

Reaction time: 5h

Reaction temperature: 110°C

15 Purification: Chromatography using acetone: CH_2Cl_2 1:10

Product: Compound 118, 0.05g oil

^{13}C NMR (CDCl_3): δ 196.6, 145.4, 144.7, 139.9, 138.6, 138.4, 138.3, 134.8, 133.0, 131.6, 131.6, 131.4, 130.2, 128.3, 125.5, 120.8, 118.1, 114.3, 20.7

20 Example 19: 4-(5-Carbonitrile-3-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 119),

General procedure: 1

Starting compound II: 5-bromonicotinonitrile (0.22g, 1.2mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g,
25 1.0 mmol)

Solvent: Toluene (5mL)

Base: Cs_2CO_3

Reaction time: 18h

Reaction temperature: 100°C

30 Purification: Chromatography using EtOAc:petroleum ether 1:2

Product: Compound 119, 0.11g oil

^{13}C NMR (DMSO): δ 195.6, 145.8, 144.5, 144.5, 138.4, 138.1, 137.1, 133.0, 131.3, 131.3, 129.8, 129.5, 126.6, 125.8, 117.3, 116.9, 114.2, 109.3, 20.1

35 Example 20: 4-(3-Bromo-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 120),

General procedure: 1

Starting compound II: 3-bromo-2-chloro-pyridine (0.23g, 1.2mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g, 1.0mmol)

Solvent: 1,4-dioxane (5mL)

5 Base: *t*-BuONa

Reaction time: 18h

Reaction temperature: 100°C

Purification: reversed phase HPLC, (0.1% TFA in H₂O):(0.1% TFA in 90% CH₃CN + 10% H₂O) 1:1 - 0:1

10 Product: Compound 120

¹³C NMR (CDCl₃): δ 196.7, 150.8, 146.5, 143.3, 140.6, 138.5, 133.8, 132.1, 131.5, 131.3, 130.3, 125.4, 119.9, 117.2, 116.4, 107.0, 20.7

Example 21: 2-Chloro-2'-methyl-4-(8-quinolylamino)benzophenone (Compound 121),

15 General procedure: 1

Starting compound II: 8-bromo-quinoline (0.16mL, 1.25mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.28g, 1.14mmol)

Solvent: 1,4-dioxane (5mL)

20 Base: *t*-BuONa

Reaction time: 18h

Reaction temperature: 100°C

Purification: Flash chromatography using Et₂O : petroleum ether 1:3

Product: Compound 121, 0.28g crystalline compound which was recrystallised from Et₂O

25 ¹³C NMR (CDCl₃): δ 196.5, 147.9, 145.9, 139.0, 138.9, 138.1, 137.7, 136.4, 134.7, 133.1, 131.4, 131.0, 130.6, 129.9, 128.8, 127.0, 125.4, 121.9, 119.1, 118.7, 115.3, 110.7, 20.6

Example 22: 2-Chloro-4-(6-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 122),

30 General procedure: 1

Starting compound II: 5-bromo-2-ethoxy-pyridine (0.24g, 1.2mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g, 1.0mmol)

35 Solvent: 1,4-dioxane (5mL)

Base: *t*-BuONa

Reaction time: 18h

Reaction temperature: 100°C

Purification: reversed phase HPLC

Product: Compound 122

^1H NMR (CDCl_3): δ 8.09(d,1H), 7.54(dd,1H), 7.40-7.20(m,5H), 6.80(m,2H), 6.66(dd,1H), 4.35(q,2H), 2.43(s,3H), 1.42(t,3H)

5

Example 23: 4-(4-Bromo-1-isoquinolylamino)-2-chloro-2'-methylbenzophenone (Compound 123),

General procedure: 1

Starting compound II: 4-bromo-1-chloroisoquinoline (0.14g, 0.5mmol)

10 Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.13g, 0.56mmol)

Solvent: 1,4-dioxane (3mL)

Base: *t*-BuONa

Reaction time: 18h

15 Reaction temperature: 100°C

Purification: reversed phase HPLC

Product: Compound 123

^1H NMR (CDCl_3): δ 8.24(s,1H), 8.18(d,1H), 8.00(d,1H), 7.85(t,1H), 7.77(d,1H), 7.65(t,1H), 7.45-7.35(m,4H), 7.28(d,1H), 7.20(t,1H), 2.52(s,3H)

20

Example 24: 2-Chloro-2'-methyl-4-(2-methyl-5-trifluoromethyl-3-pyridylamino)benzophenone (Compound 124),

General procedure: 1, without $\text{Pd}_2(\text{dba})_3$ and BINAP

Starting compound II: 3-Chloro-2-methyl-5-(trifluoromethyl)pyridine (0.20, 1.0mmol)

25 Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g, 1.0mmol)

Solvent: 1,4-dioxane (5mL)

Base: *t*-BuONa

Catalyst: $\text{Pd}(\text{P}(\text{cyclohexyl})_3)_2\text{Cl}_2$ (30 mg, 0.04mmol)

30 Reaction time: 18h

Reaction temperature: 110°C

Purification: reversed phase HPLC

Product: Compound 124

35 ^{13}C NMR (CDCl_3): δ 196.4, 152.6, 144.5, 138.8, 137.9, 137.5, 137.2, 134.8, 133.0, 132.9, 131.7, 130.4, 125.6, 123.8, 119.4, 115.4, 20.8, 19.5

Example 25: 2-Chloro-2'-methyl-4-(3-quinolylamino)benzophenone (Compound 125),

General procedure: 1

Starting compound II: 3-bromo-quinoline (0.23g, 1.1mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2)(0.25g, 1.0mmol)

5 Solvent: 1,4-dioxane (5mL)

Base: *t*-BuONa

Reaction time: 18h

Reaction temperature: 100°C

Purification by flash chromatography using EtOAc:petroleum ether 2:3

10 Product: Compound 125, 0.23g amorphous compound.

¹³C NMR (CDCl₃): δ 196.5, 146.6, 146.1, 144.9, 138.7, 138.2, 135.0, 134.2, 133.3, 131.4, 131.2, 130.7, 130.0, 129.2, 128.4, 128.0, 127.5, 126.9, 125.5, 122.4, 117.2, 113.6, 20.6

15 Example 26: 2-Chloro-4-(4-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 126),

General procedure: 1

Starting compound II: 3-bromo-4-ethoxypyridine (0.22 g, 1.1mmol)

20 Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g, 1.0mmol)

Solvent: 1,4-dioxane (6mL)

Base: *t*-BuONa

Reaction time: 7h

Reaction temperature: 100°C

25 Purification by flash chromatography using EtOAc:petroleum ether 1:3 - 1:0 followed by reversed phase HPLC.

Product: Compound 126, oil.

¹³C NMR (CDCl₃): δ 196.3, 158.5, 142.6, 138.9, 137.5, 136.5, 134.3, 134.1, 132.5, 131.9, 131.8, 131.6, 130.7, 126.8, 125.7, 120.8, 116.8, 107.7, 66.9, 20.9, 14.2

30

Example 27: 2-Chloro-4-(2-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 127),

General procedure: 1

Starting compound II: 3-bromo-2-ethoxypyridine (0.22 g, 1.1mmol)

35 Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g, 1.0mmol)

Solvent: 1,4-dioxane (6mL)

Base: *t*-BuONa

Reaction time: 3h

Reaction temperature: 100°C

Purification by flash chromatography using EtOAc:petroleum ether 1:5

Product: Compound 127, 0.20g oil.

- 5 ^{13}C NMR (CDCl_3): δ 196.4, 154.1, 145.9, 138.8, 138.7, 138.2, 134.8, 133.1, 131.4, 131.1, 130.5, 129.9, 125.4, 125.2, 122.8, 117.8, 116.6, 114.3, 62.2, 20.5, 14.7

Example 28: 2-Chloro-4-((4-isoquinolyl-N-oxide)amino)-2'-methylbenzophenone (Compound 128)

- 10 General procedure: 5

Starting compound I: Compound 114 (0.37g, 1.0mmol)

Reaction time: 2h

Purification by chromatography using EtOAc:MeOH 8:1

Product: Compound 128, 0.22g crystalline compound

- 15 ^{13}C NMR (DMSO): δ 195.5, 147.5, 138.3, 137.0, 136.0, 132.9, 131.2, 131.2, 130.9, 129.9, 129.7, 129.5, 127.9, 125.7, 125.3, 123.4, 122.1, 117.7, 114.6, 20.0

Example 29: 2-Chloro-2'-methyl-4-(2-methyl-6-quinolylamino)benzophenone (Compound 129),

- 20 General procedure: 1

Starting compound II: 6-bromoquinaldine (0.24 g, 1.1mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g, 1.0mmol)

Solvent: 1,4-dioxane (6mL)

- 25 Base: *t*-BuONa

Reaction time: 18h

Reaction temperature: 100°C

Purification by flash chromatography using EtOAc:petroleum ether 1:1

Product: Compound 129, 0.27g oil.

- 30 ^{13}C NMR (CDCl_3): δ 196.6, 157.7, 147.3, 144.9, 139.0, 138.0, 137.9, 135.2, 135.0, 133.4, 131.3, 131.0, 130.2, 129.9, 129.8, 127.3, 125.4, 124.6, 122.7, 117.1, 115.2, 113.5, 25.1, 20.5

Example 30: 2-Chloro-4-(7-chloro-4-quinolylamino)-2'-methylbenzophenone (Compound 130),

- 35 4,7-Dichloroquinoline (0.20 g, 1.0mmol) was dissolved in *i*-PrOH (10 mL) and 4-amino-2-chloro-2'-methylbenzophenone (compound 2) (0.27g, 1.1mmol) was added. The reaction

mixture was stirred at reflux for 2 h, after which it was poured into saturated aqueous NaHCO_3 (30 mL). The water phase was extracted with EtOAc (3 x 30 mL), the combined organic phases were washed with brine, dried (MgSO_4), filtered and concentrated *in vacuo*. Purification by flash chromatography using EtOAc:petroleum ether 1 : 3 - 1 : 1.

5 Product: Compound 130, 0.28g crystalline compound.

^{13}C NMR (DMSO): δ 195.67, 152.08, 149.58, 145.60, 145.01, 137.84, 137.39, 134.19, 132.34, 132.26, 131.50, 131.39, 129.82, 127.78, 125.76, 125.65, 124.51, 120.52, 119.25, 117.52, 105.38, 20.13

10 Example 31: 2-Chloro-2'-methyl-4-(2-quinolylamino)benzophenone (Compound 131), 2-Chloroquinoline (0.16 g, 1.0mmol) was dissolved in i-PrOH (10 mL) and 4-amino-2-chloro-2'-methylbenzophenone (compound 2) (0.27g, 1.1mmol) was added. The reaction mixture was stirred at reflux for 18 h, and the reaction mixture was worked up as described for the synthesis of Compound 130

15 Purification by flash chromatography using EtOAc:petroleum ether 1 : 4.

Product: Compound 131, 0.24g oil

^{13}C NMR (CDCl_3): δ 196.9, 152.6, 147.1, 144.1, 138.6, 138.4, 138.1, 134.0, 132.4, 131.6, 131.4, 131.2, 130.2, 130.1, 127.4, 127.3, 125.5, 124.5, 124.1, 119.8, 116.1, 112.9, 20.7

20

Example 32: 2-Chloro-2'-methyl-4-(4-quinolylamino)benzophenone (Compound 132), 4-Chloroquinoline (0.16g, 1.0mmol) was dissolved in *I*-PrOH (10 mL) and 4-amino-2-chloro-2'-methylbenzophenone (compound 2) (0.27g, 1.1mmol) was added. The reaction mixture was stirred at reflux for 18 h, after which it was poured into saturated aqueous NaHCO_3 (30mL). The water phase was extracted with EtOAc (3 x 30 mL), the combined organic phases were washed with brine, dried (MgSO_4), filtered and concentrated *in vacuo*. Purification by flash chromatography using EtOAc:petroleum ether 1 : 3 - 1 : 1.

25

Product: Compound 132, 0.16g amorphous compound.

^{13}C NMR (CDCl_3): δ 196.58, 150.79, 149.37, 145.25, 144.37, 138.74, 137.98, 134.30, 133.22, 132.51, 131.67, 131.61, 130.45, 130.28, 129.84, 126.10, 125.54, 121.10, 120.75, 120.07, 117.54, 105.43, 20.80

30

Example 33: 2-Chloro-2'-methyl-4-(1-methyl-7-indolylamino)benzophenone (Compound 133),

35 7-Bromoindole (0.37g, 1.9mmol) was dissolved in dry DMF. K_2CO_3 (0.52g, 3.8mmol) and methyl iodide (0.13mL, 2.08mmol) were added. The reaction was stirred at room temperature for 5 days, after which H_2O (20mL) was added, the water phase was

extracted with Et₂O (3 x 20mL), and the combined organic phases were washed with brine, dried (MgSO₄), filtered and concentrated in vacuo. The crude product was purified by chromatography, eluent EtOAc:petroleum ether 1:20 to give 7-bromo-1-methylindole as a crystalline compound.

5 ¹³C NMR (CDCl₃): δ 133.07, 131.75, 126.54, 120.46, 120.34, 103.89, 101.18, 36.84

Compound 133 was prepared according to

General procedure: 1

Starting compound II: 7-bromo-1-methylindole (0.23g, 1.1mmol)

10 Starting compound III: 4-amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g, 1.0mmol)

Solvent: 1,4-dioxane (6mL)

Base: *t*-BuONa

Reaction time: 18h

15 Reaction temperature: 100°C

Purification: Chromatography using EtOAc:petroleum ether 1:3

Product: compound 133, 0.23g amorphous compound.

¹³C NMR (CDCl₃): δ 196.55, 151.82, 139.37, 137.71, 135.36, 133.85, 132.97, 131.62, 131.19, 131.05, 130.68, 129.50, 127.92, 125.30, 123.76, 121.83, 120.53, 120.19, 20
20 115.06, 111.34, 101.50, 35.49, 20.35

Example 34: 2-Chloro-2'-methyl-4-(1-methyl-5-indolylamino)benzophenone (Compound 134),

5-Bromo-1-methylindole was prepared from 5-bromoindole as described for the
25 preparation of 7-bromo-1-methylindole from 7-bromoindole. Purification by chromatography using EtOAc:petroleum ether 1:10

¹³C NMR (CDCl₃): δ 135.37, 130.12, 129.94, 124.30, 123.27, 112.66, 110.63, 100.53, 32.96

30 Compound 134 was prepared according to

General procedure: 1

Starting compound II: 5-bromo-1-methylindole (0.23g, 1.1mmol)

Starting compound III: 4-amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25g, 1.0mmol)

35 Solvent: 1,4-dioxane (6mL)

Base: *t*-BuONa

Reaction time: 18h

Reaction temperature: 100°C

Purification: Chromatography using EtOAc:petroleum ether 1:3

Product: compound 134, 0.12g crystalline compound.

^{13}C NMR (CDCl_3): δ 7.45(d,1H), 7.36-7.18(m,6H), 7.08(d,1H), 7.06(dd,1H), 6.82(d,1H), 6.66(dd,1H), 6.45(dd,1H), 6.01(s,1H), 3.80(s,3H), 2.41(s,3H)

5

Example 35: 2-Chloro-2',5'-dimethyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 135)

General procedure: 1

Starting compound II: 3-Bromo-4-methylpyridine (0.092 g, 0.5mmol)

10 Starting compound III: 4-Amino-2-Chloro-2',5'-dimethylbenzophenone (0.134 g, 0.5mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 6 h

15 Reaction temperature: 100°C

Purification: Flash chromatography using CH_2Cl_2 :acetone 9:1

Product: Compound 135, 0.078g oil.

^{13}C NMR (CDCl_3): δ 196.7, 148.4, 146.5, 146.0, 142.1, 138.9, 135.5, 135.1, 134.9, 134.8, 133.5, 131.7, 131.2, 130.1, 129.4, 125.9, 116.0, 112.3, 20.8, 19.9, 17.5

20

Example 36: 2-Chloro-2',5'-dimethyl-4-(4-isoquinolylamino)benzophenone (Compound 136)

General procedure: 1

Starting compound II: 4-Bromoisoquinoline (0.103g, 0.49mmol)

25 Starting compound III: 4-Amino-2-Chloro-2',5'-dimethylbenzophenone (0.131g, 0.5mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 4h

30 Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc : petroleum ether 1:1

Product: Compound 136, 0.18g crystalline compound.

^{13}C NMR (CDCl_3): δ 196.8, 150.1, 149.2, 139.0, 138.6, 135.1, 134.9, 134.8, 133.5, 132.1, 131.7, 131.2, 131.1, 130.8, 130.1, 129.4, 129.3, 128.2, 127.9, 121.6, 116.4, 112.6, 20.8, 20.0

35

Example 37 2-Chloro-4-(4-Isoquinolylamino)-2',4',5'-trimethylbenzophenone (Compound 137)

General procedure: 1

Starting compound II: 4-Bromoisoquinoline (0.103 g, 0.49mmol)

- 5 Starting compound III: 4-Amino-2-chloro-2',4',5'-trimethylbenzophenone (0.148g, 0.5mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 6h

- 10 Reaction temperature: 100°C

Purification: Flash chromatography using CH₂Cl₂:acetone 9:1

Product: Compound 137, 0.16g crystalline compound.

- ¹³C NMR (CDCl₃): δ 196.6, 149.9, 148.9, 140.3, 138.3, 136.2, 135.8, 134.6, 133.5, 133.0, 132.8, 132.0, 131.5, 131.3, 130.7, 130.0, 129.3, 128.1, 127.8, 121.6, 116.4, 112.7, 20.1, 19.7, 19.1
- 15

Example 38: 2,3'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 138)

General procedure: 1

- 20 Starting compound II: 3-Bromo-4-methylpyridine (0.095 g, 0.5mmol)

Starting compound III: 4-Amino-2,3'-dichloro-2'-methylbenzophenone (0.144 g, 0.5mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

- 25 Reaction time: 6 h

Reaction temperature: 100°C

Purification: Flash chromatography using CH₂Cl₂:acetone 9:1

Product: Compound 138, 0.09g oil.

- ¹³C NMR (CDCl₃): δ 195.3, 149.2, 146.8, 146.4, 142.5, 142.0, 135.9, 135.8, 135.1, 134.9, 134.3, 131.2, 127.9, 126.9, 126.4, 126.0, 116.0, 112.1, 17.5, 17.1
- 30

Example 39: 2-Fluoro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 139)

General procedure: 1

- 35 Starting compound II: 3-Bromo-4-methylpyridine (0.092 g, 0.5mmol)

Starting compound III: 4-Amino-2-fluoro-2'-methylbenzophenone (0.120 g, 0.5mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 6 h

Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc : petroleum ether 1:1

Product: Compound 139, 0.02g oil.

- 5 ^{13}C NMR (CDCl_3): δ 194.3, 163.7, 151.3, 146.8, 146.6, 142.8, 140.3, 136.3, 135.2, 134.0, 130.9, 130.2, 128.1, 126.0, 125.3, 117.9, 109.9, 101.1, 19.9, 17.4

Example 40: 2,4'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone
(Compound 140)

- 10 General procedure: 1

Starting compound II: 3-Bromo-4-methylpyridine (0.086 g, 0.5mmol)

Starting compound III: 4-Amino-2,4'-dichloro-2'-methylbenzophenone (0.132g, 0.47mmol)

Solvent: 1,4-dioxan (5mL)

- 15 Base: *t*-BuONa

Reaction time: 6 h

Reaction temperature: 100°C

Purification: Flash chromatography using CH_2Cl_2 :acetone 9:1

Product: Compound 140, 0.12g oil.

- 20 ^{13}C NMR (CDCl_3): δ 195.4, 148.7, 146.7, 146.2, 142.2, 140.1, 137.4, 136.8, 135.3, 135.2, 133.4, 131.3, 131.1, 128.9, 125.9, 125.6, 115.8, 112.4, 20.4, 17.5

Example 41: 2-Chloro-4'-fluoro-4-(4-isoquinolylamino)-2'-methylbenzophenone
(Compound 141)

- 25 General procedure: 1

Starting compound II: 4-Bromoisoquinoline (0.104g, 0.5mmol)

Starting compound III: 4-Amino-2-chloro-4'-fluoro-2'-methylbenzophenone (0.140g, 0.53mmol)

Solvent: 1,4-dioxan (5mL)

- 30 Base: *t*-BuONa

Reaction time: 6 h

Reaction temperature: 100°C

Purification: Flash chromatography using CH_2Cl_2 :acetone 9:1

Product: Compound 141, 0.13g crystalline compound.

- 35 ^{13}C NMR (CDCl_3): δ 195.4, 164.0, 150.3, 149.3, 141.8, 138.7, 135.0, 134.9, 133.1, 132.4, 132.4, 131.0, 130.8, 129.4, 129.3, 128.2, 127.9, 121.6, 118.2, 116.2, 112.7, 112.3, 20.8

Example 42: 4'-*n*-Butyl-2-chloro-4-(4-isoquinolylamino)-2'-methylbenzophenone
(Compound 142)

General procedure: 1

- 5 Starting compound II: 4-Bromoisquinoline (0.104g, 0.5mmol)
Starting compound III: 4-Amino-2'-*n*-butyl-2-chloro-2'-methylbenzophenone (0.150g, 0.5mmol)
Solvent: 1,4-dioxan (5mL)
Base: *t*-BuONa
- 10 Reaction time: 6 h
Reaction temperature: 100°C
Purification: Flash chromatography using CH₂Cl₂:acetone 9:1
Product: Compound 142, 0.17g oil.
¹³C NMR (CDCl₃): δ 196.4, 150.0, 148.8, 146.7, 138.6, 138.5, 136.0, 134.7, 133.0,
15 132.1, 131.7, 131.2, 130.8, 130.7, 130.1, 129.3, 128.2, 127.9, 125.4, 121.6, 116.3,
112.7, 35.6, 33.3, 22.4, 20.8, 13.9

Example 43: 2-Chloro-4-(5-isoquinolylamino)-2'-methylbenzophenone (Compound 143)

General procedure: 1

- 20 Starting compound II: 5-Bromoisquinoline (0.104g, 0.5mmol)
Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.122g, 0.5mmol)
Solvent: 1,4-dioxan (5mL)
Base: *t*-BuONa
- 25 Reaction time: 6 h
Reaction temperature: 100°C
Purification: Flash chromatography using CH₂Cl₂:acetone 9:1
Product: Compound 143, 0.16g oil.
¹³C NMR (CDCl₃): δ 196.6, 153.0, 148.7, 143.4, 138.9, 138.0, 135.5, 135.0, 133.5,
30 131.6, 131.3, 131.0, 129.8, 129.5, 127.4, 125.4, 124.8, 123.8, 116.6, 115.1, 113.0, 20.5

Example 44: 2-Chloro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone
(Compound 144)

General procedure: 1

- 35 Starting compound II: 4-Bromoisquinoline (0.104g, 0.5mmol)
Starting compound III: 4-Amino-2-chloro-4'-methoxy-2'-methylbenzophenone (0.136g, 0.5mmol)
Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 6 h

Reaction temperature: 100°C

Purification: Flash chromatography using CH₂Cl₂:acetone 9:1

5 Product: Compound 144, 0.19g crystalline compound.

¹³C NMR (CDCl₃): δ 195.5, 161.9, 149.9, 148.5, 142.0, 138.3, 134.2, 133.7, 132.4, 132.0, 131.4, 130.9, 130.7, 129.3, 128.2, 127.8, 121.6, 117.1, 116.3, 112.9, 110.4, 55.3, 21.5

10 Example 45: 2-Fluoro-4-(4-Isoquinolylamino)-4'-methoxy-2'-methylbenzophenone (Compound 145)

General procedure: 1

Starting compound II: 4-Bromoisoquinoline (0.103g, 0.5mmol)

15 Starting compound III: 4-Amino-2-fluoro-4'-methoxy-2'-methylbenzophenone (0.130g, 0.5mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 6 h

Reaction temperature: 100°C

20 Purification: Flash chromatography using EtOAc : petroleum ether 1:1 followed by recrystallization from EtOAc.

Product: Compound 145, 0.07g crystalline compound.

25 ¹H NMR (DMSO-D₆): δ 9.18 (s,1H), 9.13 (s,1H), 8.54 (s,1H), 8.20 (d,1H), 8.03 (d,1H), 7.83 (t,1H), 7.75 (t,1H), 7.42 (t,1H), 7.30 (d,1H), 6.87 (d,1H), 6.81 (dd,1H), 6.77 (dd,1H), 6.58 (dd,1H), 3.80 (s,3H), 2.34 (s,3H)

Example 46: 2,4'-Dichloro-2'-methyl-4-(1-methyl-7-Indolylamino)benzophenone (Compound 146)

General procedure: 1

30 Starting compound II: 7-Bromo-1-methylindole (0.115g, 0.55mmol) prepared as described in example 33

Starting compound III: 4-Amino-2,4'-dichloro-2'-methylbenzophenone (0.140g, 0.5mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

35 Reaction time: 6 h

Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc : petroleum ether 1:3

Product: Compound 146, 0.08g crystalline compound.

^{13}C NMR (CDCl_3): δ 195.5, 152.1, 139.8, 137.8, 136.5, 135.4, 133.8, 132.9, 131.6, 131.2, 131.1, 130.8, 127.5, 125.5, 123.6, 121.8, 120.6, 120.2, 115.0, 111.4, 101.5, 35.5, 20.2

5 **Example 47:** 2-Chloro-4-(1-methyl-7-indolylamino)-2',4',5'-trimethylbenzophenone (Compound 147)

General procedure: 1

Starting compound II: 7-Bromo-1-methylindole (0.115g, 0.55mmol) prepared as described in example 33

10 Starting compound III: 4-Amino-2-chloro-2',4',5'-trimethylbenzophenone (0.137g, 0.5mmol)

Solvent: 1,4-dioxan (5 mL)

Base: *t*-BuONa

Reaction time: 6 h

15 Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc : petroleum ether 1:3

Product: Compound 147, 0.097g crystalline compound.

^{13}C NMR (CDCl_3): δ 196.6, 151.5, 140.0, 136.6, 135.5, 135.0, 133.5, 133.4, 133.0, 132.7, 131.6, 131.3, 131.0, 128.6, 124.0, 121.8, 120.4, 120.2, 115.0, 111.3, 101.5, 35.5, 20.0, 19.7, 19.1

Example 48: 2-Chloro-2',5'-dimethyl-4-(1-methyl-7-indolylamino)benzophenone (Compound 148)

General procedure: 1

25 Starting compound II: 7-Bromo-1-methylindole (0.115g, 0.55mmol) prepared as described in example 33

Starting compound III: 4-Amino-2-chloro-2',5'-dimethylbenzophenone (0.148 g, 0.5mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

30 Reaction time: 6 h

Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc : petroleum ether 1:3

Product: Compound 148, 0.30g crystalline compound.

^1H NMR (CDCl_3): δ 7.56 (d,1H), 7.28 (d,1H), 7.17-7.02 (m,4H), 6.98 (d,1H), 6.95 (d,1H), 6.61 (d,1H), 6.50 (d,1H), 6.42 (dd,1H), 5.93 (s,Br,1H), 3.81 (s,3H), 2.32 (s,3H), 2.27 (s,3H)

Example 49: 2-Chloro-4-(3-ethoxy-4-isoquinolylamino)-2'-methylbenzophenone

(Compound 149)

General procedure: 1

Starting compound II: 4-Bromo-3-ethoxyisoquinoline (0.097g, 0.39mmol)

5 Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.095g, 0.39mmol)

Solvent: 1,4-dioxan (3mL)

Base: t-BuONa

Reaction time: 6 h

10 Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc : petroleum ether 1:3

Product: Compound 149, 0.11g crystalline compound.

¹³C NMR (CDCl₃): δ 196.6, 155.1, 149.9, 147.4, 139.3, 137.8, 134.8, 133.3, 131.2,
130.7, 130.5, 129.6, 128.7, 128.1, 125.6, 125.3, 124.5, 121.9, 116.2, 113.9, 112.4,
15 62.7, 20.4, 15.0

Example 50: 2-Chloro-4-(1-ethoxy-4-isoquinolylamino)-2'-methylbenzophenone

(Compound 150)

General procedure: 1

20 Starting compound II: 4-Bromo-1-ethoxyisoquinoline (0.097g, 0.39mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.095g, 0.39mmol)

Solvent: 1,4-dioxan (3mL)

Base: t-BuONa

25 Reaction time: 6 h

Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc : petroleum ether 1:3

Product: Compound 150, 0.11g crystalline compound.

¹³C NMR (CDCl₃): δ 196.5, 159.7, 151.2, 139.4, 138.5, 137.7, 135.5, 135.3, 133.8,
30 131.2, 131.0, 130.7, 129.5, 128.0, 127.1, 125.3, 125.0, 124.6, 121.8, 120.3, 115.1,
111.4, 62.4, 20.4, 14.6

Example 51: 4-(2-Benzoxazolylamino)-2-chloro-2'-methylbenzophenone (Compound 151)

General procedure: 1

35 Starting compound II: 2-Chlorobenzoxazole (0.35mL, 3.0mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.49g, 2.0mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 16 h

Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc : petroleum ether 1:4

5 Product: Compound 151, 0.17g, crystalline compound.

¹³C NMR (CDCl₃): δ 196.9, 157.3, 147.8, 141.4, 141.3, 138.7, 138.0, 134.0, 133.1, 132.2, 131.6, 130.6, 125.5, 124.7, 124.5, 122.6, 119.3, 117.4, 115.7, 109.5, 20.8

Example 52 4-(1-Methyl-2-benzimidazolylamino)-2-chloro-2'-methylbenzophenone

10 (Compound 152)

General procedure: 1

Starting compound II: 2-Chloro-1-methylbenzimidazole (0.33g, 2.0mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.49 g, 2.0mmol)

15 Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 24 h

Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc : petroleum ether 1 : 2

20 Product: Compound 152, 0.27g, crystalline compound

¹³C NMR (CDCl₃): δ 196.0, 147.0, 140.4, 139.0, 137.1, 135.4, 133.9, 132.0, 131.8, 130.9, 129.8, 125.6, 125.4, 124.6, 121.1, 117.6, 113.7, 110.1, 31.3, 20.9

Example 53 4-(2-Benzothiazolylamino)-2-chloro-2'-methylbenzophenone (Compound

25 153)

General procedure: 1

Starting compound II: 2-Chlorobenzothiazole (0.26mL, 2.0mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.49g, 2.0mmol)

30 Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 20 h

Reaction temperature: 100°C

Purification: Flash chromatography using acetone:CH₂Cl₂ 1:100

35 Product: Compound 153, 0.24g, crystalline compound

¹³C NMR (CDCl₃): δ 196.9, 161.5, 151.1, 143.1, 138.7, 138.0, 134.1, 133.1, 132.3, 131.6, 130.5, 130.3, 126.5, 125.5, 123.4, 121.0, 120.3, 119.6, 116.0, 20.8

Example 54 2-Chloro-2'-methyl-4-(2-pyrimidylamino)benzophenone (Compound 154)

General procedure: 1

Starting compound II: 2-Bromopyrimidine (0.18g, 1.1mmol)

5 Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25 g, 1.0mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 18 h

10 Reaction temperature: 100°C

Purification: Flash chromatography using acetone:CH₂Cl₂ 2:100

Product: Compound 154, 0.14g, crystalline compound

¹³C NMR (CDCl₃): δ 196.8, 159.4, 158.1, 143.0, 138.4, 133.9, 132.1, 132.0, 131.5, 131.3, 130.3, 125.4, 119.8, 116.2, 113.8, 20.7

15

Example 55 2-Chloro-2'-methyl-4-(7-methyl-purin-6-ylamino)benzophenone (Compound 155)

General procedure: 1

Starting compound II: 6-Bromo-7-methylpurine (0.18g, 0.84mmol)

20 Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.19g, 0.77mmol)

Solvent: 1,4-dioxan (5mL)

Base: *t*-BuONa

Reaction time: 18 h

25 Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc:petroleumether 1 : 3 -> 1 : 0 followed by recrystallisation from Et₂O

Product: Compound 155, 0.06g crystalline compound

30 ¹³C NMR (CDCl₃): δ 196.7, 152.7, 151.4, 150.4, 142.3, 141.8, 138.7, 138.2, 133.7, 133.1, 131.9, 131.6, 131.4, 130.5, 125.5, 120.7, 120.6, 117.1, 30.0, 20.8

Example 56 2-Chloro-2'-methyl-4-(2-methyl-5-benzothiazolylamino)benzophenone (Compound 156)

General procedure: 1

35 Starting compound II: 5-Bromo-2-methylbenzothiazole (0.25g, 1.1mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25 g, 1.0mmol)

Solvent: 1,4-dioxan (6mL)

Base: *t*-BuONa

Reaction time: 18 h

Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc:petroleumether 1 : 3 -> 1 : 2

5 Product: Compound 156, 0.21g oil

¹³C NMR (CDCl₃): δ 196.5, 168.7, 154.6, 147.8, 139.1, 138.8, 138.0, 135.1, 133.4, 131.3, 130.9, 130.8, 129.8, 129.5, 125.4, 122.2, 119.3, 116.5, 114.3, 113.0, 20.5, 20.2

Example 57 2-Chloro-2'-methyl-4-(pyrazin-2-ylamino)benzophenone (Compound 157)

10 General procedure: 1

Starting compound II: 2-Iodopyrazine (0.23g, 1.1mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25 g, 1.0mmol)

Solvent: 1,4-dioxan (6mL)

15 Base: *t*-BuONa

Reaction time: 6 h

Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc:petroleum ether 1 : 1 followed by purification on RP HPLC

20 Product: Compound 157, oil

¹³C NMR (CDCl₃): δ 196.8, 151.0, 143.2, 141.6, 138.6, 138.3, 136.3, 134.5, 134.0, 132.4, 132.2, 131.6, 131.4, 130.4, 125.5, 119.7, 116.1, 20.7

Example 58 2-Chloro-2'-methyl-4-(5-pyrimidylamino)benzophenone (Compound 158)

25 General procedure: 1

Starting compound II: 5-Bromopyrimidine (0.18g, 1.1mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.25 g, 1.0mmol)

Solvent: 1,4-dioxan (5mL)

30 Base: *t*-BuONa

Reaction time: 6 h

Reaction temperature: 100°C

Purification: Flash chromatography using EtOAc:petroleum ether 1:1 -> 1:0 followed by purification on RP HPLC

35 Product: Compound 158, oil

¹³C NMR (CDCl₃): δ196.3, 153.3, 148.0, 145.0, 138.6, 138.2, 134.9, 133.0, 132.2, 131.6, 131.5, 130.2, 125.5, 117.7, 114.2, 20.7

Example 59 2-Chloro-2'-methyl-4-(5-nitro-2-thiazolylamino)benzophenone (Compound 159)

4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.246g, 1.0mmol), 2-bromo-5-nitrothiazol (0.209g, 1.0mmol) and N,N-diisopropyl ethylamine (0.18mL, 1.0mmol) were mixed in a 4 mL screwcap vessel and kept at 60°C for 18h. The reaction mixture was partitioned between EtOAc and water, and the organic phase was washed with H₂O (3 x 10 mL) and brine (10mL), dried (MgSO₄) and evaporated *in vacuo*. The crude compound was purified by flash chromatography using EtOAc:CH₂Cl₂ 1:50 as the eluent.

Product: Compound 159, 0.12g, crystalline compound.

¹³C NMR (DMSO-D-6): δ 195.8, 166.5, 145.2, 142.1, 138.6, 137.8, 137.4, 133.1, 131.9, 131.8, 131.6, 130.3, 125.9, 119.3, 117.0, 54.9, 20.3

Example 60 2-Chloro-2'-methyl-4-((4-methyl-3-nitro-(1,2,4-triazol-5-ylamino))benzophenone (Compound 160)

Preparation of 5-bromo-4-methyl-3-nitro-1,2,4-triazole:

5-Bromo-3-nitro-1,2,4-triazole (0.78g, 4.0 mmol) was dissolved in DMF (20mL). K₂CO₃ (1.11g, 8.0mmol) and CH₃I (0.28mL, 4.4mmol) were added, and the mixture was stirred at room temperature for 18h. H₂O was added, and the water phase was extracted three times with Et₂O. The combined organic phases were washed with brine, dried (MgSO₄) and evaporated, and the crude product was purified by flash chromatography using EtOAc:petroleum ether 1:1 as the eluent

Product: 2-bromo-1-methyl-5-nitro-1,2,4-triazole, 0.44g

¹³C NMR (CDCl₃): δ 131.3, 80.6, 38.0

Preparation of compound 160:

General procedure: 1

Starting compound II: 5-Bromo-4-methyl-3-nitro-1,2,4-triazole (0.22g, 1.1mmol)

Starting compound III: 4-Amino-2-chloro-2'-methylbenzophenone (compound 2) (0.24 g, 1.0mmol)

Solvent: Toluene (5mL)

Base: Cs₂CO₃

Reaction time: 18 h

Reaction temperature: 100°C

After 18h an extra portion of BINAP (24mg) and Pd₂(dba)₃ (17mg) were added, and the reaction mixture was shaken another 24h at 100°C.

Work up as described in General procedure 1

Purification: Flash chromatography using EtOAc:petroleum ether 1 : 1 followed by purification on reversed phase HPLC

Product: Compound 160, oil

¹³C NMR (DMSO-D-6): δ 195.8, 158.3, 151.0, 142.8, 137.8, 137.4, 131.9,

5 131.6, 131.4, 131.4, 129.9, 125.8, 117.9, 115.6, 35.6, 20.1

Example 61

Tablet containing compound 116

	Compound 116 (active substance)	50 mg
10	Lactose	125 mg
	Starch	12 mg
	Methyl cellulose	2 mg
	Sodium carboxymethyl cellulose	10 mg
	Magnesium stearate	1 mg

15

The active substance, lactose and starch are mixed to a homogeneous state in a suitable mixer and moistened with a 5 per cent aqueous solution of methyl cellulose 15 cps. The mixing is continued until granules are formed. If necessary, the wet granulation is passed through a suitable screen and dried to a water content of less than 1% in a suitable drier, e.g. fluid bed or drying oven. The dried granules are passed through a 1 mm screen and mixed to a homogeneous state with sodium carboxymethyl cellulose. Magnesium stearate is added, and the mixing is continued for a short period of time. Tablets with a weight of 200 mg are produced from the granulation by means of a suitable tableting machine.

20

25 Example 62: Formulation for injection containing compound 116.

	Compound 116 (active substance)	1%
	Sodium chloride	q.s.
	Ethanol	10%
	Water for injection to make	100%

30

The active substance is dissolved in ethanol (10%) then water for injection made isotonic with sodium chloride is added to make 100%. The mixture is filled into ampoules and sterilised.

35

Example 63 Cream formulation containing compound 116

Compound 116 (10 g) was dissolved in Octyldodecyl myristate (250g) to form Part A.

Methylparaben (1 g) and propylparaben (0.2 g) were dissolved in phenoxyethanol (6 g)

and mixed with a 0.025 M Phosphate buffer pH = 7.5 (632,8 g) to form Part B. Cetostearyl alcohol (50 g) and ARLACEL 165® (50 g) was melted in a vessel at 70° to 80 °C. Part A was added and heated to 60-70°C. The aqueous phase was likewise heated to 60-70 °C and slowly added to the melted oil phase under high speed stirring. The homogenised components were cooled to room temperature.

10

15

20

25


30

35

CLAIMS

1. A compound of the general formula I below

(I)



wherein

R_1 is selected from the group consisting of halogen, haloalkyl, hydroxy, hydroxyalkyl, hydroxyalkyloxy, mercapto, cyano, carboxy, nitro, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, aryl, heteroaryl, aralkyl, alkylaryl, alkoxy, aralkoxy, alkylthio, alkoxycarbonyl, alkylcarbonyloxy, alkoxycarbonyloxy, alkylsulfonyloxy, alkylloxysulfonyl, alkylcarbonylamino, aminocarboaminoalkyl, aminosulfonyl, alkylsulfonylamino, alkanoyl, alkylcarbonyl, $-NR_9R_{10}$ or $-CONR_9R_{10}$, wherein R_9 and R_{10} are the same or different and individually represent hydrogen, alkyl or aryl;

R_2 represents one or more, same or different substituents selected from the group consisting of hydrogen, halogen, haloalkyl, hydroxy, hydroxyalkyl, hydroxyalkyloxy, mercapto, cyano, carboxy, nitro, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, aryl, heteroaryl, aralkyl, alkylaryl, alkoxy, aralkoxy, alkylthio, alkoxycarbonyl, alkylcarbonyloxy, alkoxycarbonyloxy, alkylsulfonyloxy, alkylloxysulfonyl, alkylcarbonylamino, aminocarboaminoalkyl, aminosulfonyl, alkylsulfonylamino, alkanoyl, alkylcarbonyl, $-NR_9R_{10}$ or $-CONR_9R_{10}$, wherein R_9 and R_{10} are the same or different and individually represent hydrogen, alkyl or aryl;

R_3 represents one or more, same or different substituents selected from the group consisting of hydrogen, halogen, haloalkyl, hydroxy, hydroxyalkyl, mercapto, cyano, nitro, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, aralkyl, alkylaryl, alkoxy, aralkoxy, alkylthio, alkoxycarbonyl, alkylcarbonylamino, alkylcarbonyloxy, alkoxycarbonyloxy, alkylcarbonyl $-NR_9R_{10}$ or $-CONR_9R_{10}$, wherein R_9 and R_{10} are the same or different and individually represent hydrogen, alkyl or aryl;

R₄ represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, carboxy or aryl;

R₅ represents a heteroaromatic mono- or bicyclic ring system comprising 1-4 heteroatoms, except for triazine, the ring system being optionally substituted by
 5 hydrogen, halogen, haloalkyl, hydroxy, hydroxyalkyl, hydroxyalkyloxy, mercapto, cyano, carboxy, nitro, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, aryl, heteroaryl, aralkyl, alkylaryl, alkoxy, aralkoxy, alkylthio, alkoxycarbonyl, alkylcarbonyloxy, alkoxycarbonyloxy, alkylsulfonyloxy, alkylsulfonyl, alkylcarbonylamino, aminocarboaminoalkyl, aminosulfonyl, alkylsulfonylamino, alkanoyl, alkylcarbonyl, -NR₉R₁₀
 10 or -CONR₉R₁₀, wherein R₉ and R₁₀ are the same or different and individually represent hydrogen, alkyl or aryl;

X represents oxygen, sulphur, N-OH or NR₁₁ wherein R₁₁ is hydrogen or alkyl;

15 and salts thereof with pharmaceutically acceptable acids, hydrates and solvates, and optionally the N-oxides thereof wherein a nitrogen atom of the heterocyclic R₅ substituent is oxidised.

2. A compound according to claim 1 wherein R₁ represents a substituent selected from
 20 the group consisting of halogen, hydroxy, mercapto, trifluoromethyl, amino, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, (C₁-C₆)alkylamino, (C₁-C₆)alkoxycarbonyl, cyano, -CONH₂, phenyl, and nitro; in particular fluoro, chloro, bromo, hydroxy, trifluoromethyl, amino, (C₁-C₃)alkyl, (C₂-C₃)alkenyl, (C₁-C₃)alkoxy, (C₁-C₃)alkoxycarbonyl, cyano, and -CONH₂.

25 3. A compound according to claim 1 or 2 wherein R₂ represents one or more, same or different substituents selected from the group consisting of hydrogen, halogen, hydroxy, mercapto, trifluoromethyl, amino, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio, (C₁-C₆)alkylamino, (C₁-C₆)alkoxycarbonyl, cyano, -CONH₂, phenyl, and
 30 nitro; In particular hydrogen, fluoro, chloro, bromo, hydroxy, trifluoromethyl, amino, (C₁-C₃)alkyl, (C₂-C₃)alkenyl, (C₁-C₃)alkoxy, (C₁-C₃)alkoxycarbonyl, cyano, and -CONH₂.

4. A compound according to any one of claims 1-3 wherein R₃ represents one or more, same or different substituents selected from the group consisting of hydrogen, halogen,
 35 hydroxy, mercapto, trifluoromethyl, amino, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy,

(C₁-C₆)alkylthio, (C₁-C₆)alkylamino, (C₁-C₆)alkoxycarbonyl, cyano, -CONH₂, phenyl, and nitro; In particular hydrogen, fluoro, chloro, bromo, hydroxy, trifluoromethyl, amino, (C₁-C₃)alkyl, (C₂-C₃)alkenyl, and (C₁-C₃)alkoxy.


5 5. A compound according to any one of claims 1-4 wherein R₄ represents hydrogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, or (C₃-C₆) cycloalkyl or -cycloalkenyl; In particular hydrogen, (C₁-C₄)alkyl, or (C₂-C₄)alkenyl.

10 6. A compound according to any one of claims 1-5 wherein R₅ represents an optionally substituted heteroaromatic ring system having one or 2 fused rings of 5 or 6 ring atoms and comprising 1 or 2 nitrogen atoms.



7. A compound according to claim 6 wherein R₅ is selected from the group consisting of

15



20





25



30



and



wherein R_6 and R_7 represent one or more, same or different substituents selected from the group consisting of hydrogen, halogen, hydroxy, mercapto, trifluoromethyl, cyano, carboxy, carbamoyl, amino, nitro, (C_1-C_{10}) alkyl, (C_2-C_{10}) alkenyl, (C_3-C_8) cycloalkyl or -cycloalkenyl, (C_1-C_{10}) alkoxy, (C_1-C_{10}) alkylthio, (C_1-C_{10}) alkoxycarbonyl, and phenyl; and R_8 represents hydrogen, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, or (C_3-C_6) cycloalkyl or -cycloalkenyl; and wherein R_5 is optionally oxidised to the corresponding N-oxide as exemplified herein.

15

8. A compound according to any one of claims 1-7 wherein X represents oxygen, sulphur or NH, in particular oxygen or NH.

20

9. A compound according to claim 7, wherein R_6 and R_7 represent one or more, same or different substituents selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, trifluoromethyl, amino, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, (C_1-C_6) alkoxy, (C_1-C_6) alkoxycarbonyl, cyano, carboxy, and $-CONH_2$.

25


10. A compound according to any one of claims 7-9 wherein R_8 represents hydrogen, (C_1-C_4) alkyl, (C_2-C_6) alkenyl, such as allyl.

11. A compound according to claim 10 wherein R_8 represents hydrogen, methyl, ethyl, allyl, propyl, benzyl, or t-butyl.

30

12. A compound according to any one of claims 1-11 wherein R_1 represents a substituent selected from the group consisting of fluoro, chloro, bromo, hydroxy, methyl, and methoxy.

13. A compound according to any one of claims 1-12 wherein R_2 represents one or more, same or different substituents selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, trifluoromethyl, methyl, ethyl, and methoxy.
- 5 14. A compound according to any one of claims 1-13 wherein R_3 represents one or more, same or different substituents selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, methyl, and methoxy.
- 10 15. A compound according to any one of claims 1-14 wherein R_4 represents hydrogen, methyl, or ethyl.
- 15 16. A compound according to any one of claims 1-15 wherein R_5 is selected from the group consisting of substituted or non-substituted 3-pyridyl, 2-pyridyl, 3-quinolyl, 4-isoquinolyl, 4-indolyl, 5-indolyl, 6-indolyl or 7-indolyl moieties, and the corresponding N-oxides;
- 20 17. A compound according to claim 7 wherein R_6 and R_7 represents one or more, same or different substituents selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, methyl, methoxy, cyano, and carboxy.
- 25 18. A compound according to any one of claims 1-17 wherein X represents oxygen.
19. A compound according to any one of claims 1-18 characterised in that at least one of R_1 , R_2 , R_3 , R_6 and R_7 represents a phenyl group optionally substituted by hydroxy, amino, nitro, cyano, halogen, methyl or methoxy.
- 30 20. A compound according to any one of claims 1-19 wherein said halogen is selected from the group consisting of fluoro, chloro, and bromo.
- 35 21. A compound of the general formula Ic:



wherein R₁, R₂, R₄, R₅, R₆ and X have the meaning specified in formula I, and R₃ represents (C₁-C₃)alkyl, fluoro, chloro, bromo, methoxy, and hydroxy, and salts thereof with pharmaceutically acceptable acids, hydrates and solvates.

- 5 22. A compound of the general formula Ic, in which R₁ preferably represents methyl or halogen, and more preferably F or Cl; R₂ represents one or more substituents, preferably hydrogen, (C₁-C₃)alkyl, methoxy or ethoxy, and R₃ represents methyl, methoxy or chloro.

23. A compound according to claim 22 wherein R₃ represents a halogen atom.

- 10 24. A compound according to claim 23 wherein R₃ represents chlorine.

25. A compound according to formula I selected from the group consisting of:
 2-Chloro-2'-methyl-4-(4-pyridylamino)benzophenone (Compound 101),
 15 2-Chloro-2'-methyl-4-(2-pyridylamino)benzophenone (Compound 102),
 2-Chloro-2'-methyl-4-(5-nitro-2-pyridylamino)benzophenone (Compound 103),
 4-(6-Amino-5-nitro-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 104),
 6-Chloro-2-(3-chloro-4-(2-methylbenzoyl)phenylamino) isonicotinic acid (Compound 105),
 4-(6-carbonitrile-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 106),
 20 2-Chloro-2'-methyl-4-(3-pyridylamino)benzophenone (Compound 107),
 2-Chloro-4-(5,6-diamino-2-pyridylamino)-2'-methylbenzophenone (Compound 108),
 2-Chloro-2'-methyl-4-(3-nitro-2-pyridylamino)benzophenone (Compound 109),
 4-(3-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 110),
 4-(5-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 111),
 25 2-Chloro-4-(4-isoquinolylamino)-2'-methylbenzophenone (Compound 114),
 t-Butyl 5-(3-chloro-4-(2-methylbenzoyl)phenylamino) nicotinoate (Compound 115),
 2-Chloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 116),
 2-Chloro-2'-methyl-4-(6-methyl-3-pyridylamino)benzophenone (Compound 117),
 4-(5-Bromo-3-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 118),
 30 4-(5-Carbonitrile-3-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 119),
 4-(3-Bromo-2-pyridylamino)-2-chloro-2'-methylbenzophenone (Compound 120),
 2-Chloro-2'-methyl-4-(8-quinolylamino)benzophenone (Compound 121),
 2-Chloro-4-(6-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 122),
 4-(4-Bromo-1-isoquinolylamino)-2-chloro-2'-methylbenzophenone (Compound 123),
 35 2-Chloro-2'-methyl-4-(2-methyl-5-trifluoromethyl-3-pyridylamino)benzophenone
 (Compound 124),
 2-Chloro-2'-methyl-4-(3-quinolylamino)benzophenone (Compound 125),
 2-Chloro-4-(4-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 126),

- 2-Chloro-4-(2-ethoxy-3-pyridylamino)-2'-methylbenzophenone (Compound 127),
 2-Chloro-2'-methyl-4-(2-methyl-6-quinolylamino)benzophenone (Compound 129),
 2-Chloro-2'-methyl-4-(2-quinolylamino)benzophenone (Compound 131),
 2-Chloro-4-(7-chloro-4-quinolylamino)-2'-methylbenzophenone (Compound 130),
 5 2-Chloro-2'-methyl-4-(4-quinolylamino)benzophenone (Compound 132),
 2-Chloro-2'-methyl-4-(1-methyl-7-indolylamino)benzophenone (Compound 133),
 2-Chloro-2'-methyl-4-(1-methyl-5-indolylamino)benzophenone (Compound 134),
 2-Chloro-2',5'-dimethyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 135),
 2-Chloro-2',5'-dimethyl-4-(4-isoquinolylamino)benzophenone (Compound 136),
 10 2-Chloro-4-(4-isoquinolylamino)-2',4',5'-trimethylbenzophenone (Compound 137),
 2,3'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 138),
 2-Fluoro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 139),
 2,4'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone (Compound 140),
 2-Chloro-4'-fluoro-4-(4-isoquinolylamino)-2'-methylbenzophenone (Compound 141),
 15 4'-*n*-Butyl-2-chloro-4-(4-isoquinolylamino)-2'-methylbenzophenone (Compound 142),
 2-Chloro-4-(5-isoquinolylamino)-2'-methylbenzophenone (Compound 143),
 2-Chloro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone (Compound 144),
 2-Fluoro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone (Compound 145),
 2,4'-Dichloro-2'-methyl-4-(1-methyl-7-indolylamino)benzophenone (Compound 146),
 20 2-Chloro-4-(1-methyl-7-indolylamino)-2',4',5'-trimethylbenzophenone (Compound 147),
 2-Chloro-2',5'-dimethyl-4-(1-methyl-7-indolylamino)benzophenone (Compound 148),
 2-Chloro-4-(3-ethoxy-4-isoquinolylamino)-2'-methylbenzophenone (Compound 149),
 2-Chloro-4-(1-ethoxy-4-isoquinolylamino)-2'-methylbenzophenone (Compound 150),
 4-(2-Benzoxazolylamino)-2-chloro-2'-methylbenzophenone (Compound 151),
 25 4-(1-Methyl-2-benzimidazolylamino)-2-chloro-2'-methylbenzophenone (Compound 152),
 4-(2-Benzothiazolylamino)-2-chloro-2'-methylbenzophenone (Compound 153),
 2-Chloro-2'-methyl-4-(2-pyrimidylamino)benzophenone (Compound 154),
 2-Chloro-2'-methyl-4-(7-methyl-purin-6-ylamino)benzophenone (Compound 155),
 2-Chloro-2'-methyl-4-(2-methyl-5-benzothiazolylamino)benzophenone (Compound 156),
 30 2-Chloro-2'-methyl-4-(pyrazin-2-ylamino)benzophenone (Compound 157),
 2-Chloro-2'-methyl-4-(5-pyrimidylamino)benzophenone (Compound 158),
 2-Chloro-2'-methyl-4-(5-nitro-2-thiazolylamino)benzophenone (Compound 159),
 2-Chloro-2'-methyl-4-((4-methyl-3-nitro-(1,2,4-triazol-5-ylamino)))benzophenone
 (Compound 160)
 35 and the N-oxides thereof wherein the nitrogen atom of the heterocyclic R₅ substituent is
 specifically oxidised, including the N-oxides:
 2-Chloro-2'-methyl-4((2-pyridyl-N-oxide)amino)benzophenone (Compound 112),
 2-Chloro-2'-methyl-4((3-pyridyl-N-oxide)amino)benzophenone (Compound 113),

- 4-(4-Bromo-1-isoquinolylamino)-2-chloro-2'-methyl(thiobenzophenone)
 2-Chloro-2'-methyl-4-(2-methyl-5-trifluoromethyl-3-pyridylamino)thiobenzophenone
 2-Chloro-2'-methyl-4-(3-quinolylamino)thiobenzophenone
 2-Chloro-4-(4-ethoxy-3-pyridylamino)-2'-methyl(thiobenzophenone)
 5 2-Chloro-4-(2-ethoxy-3-pyridylamino)-2'-methyl(thiobenzophenone)
 2-Chloro-2'-methyl-4-(2-methyl-6-quinolylamino)thiobenzophenone
 2-Chloro-2'-methyl-4-(2-quinolylamino)thiobenzophenone
 2-Chloro-4-(7-chloro-4-quinolylamino)-2'-methyl(thiobenzophenone)
 2-Chloro-2'-methyl-4-(4-quinolylamino)thiobenzophenone
 10 2-Chloro-2'-methyl-4-(1-methyl-7-indolylamino)thiobenzophenone
 2-Chloro-2'-methyl-4-(1-methyl-5-indolylamino)thiobenzophenone
 2-Chloro-2',5'-dimethyl-4-(4-methyl-3-pyridylamino)thiobenzophenone,
 2-Chloro-2',5'-dimethyl-4-(4-isoquinolylamino)thiobenzophenone,
 2-Chloro-4-(4-isoquinolylamino)-2',4',5'-trimethyl(thiobenzophenone),
 15 2,3'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)thiobenzophenone,
 2-Fluoro-2'-methyl-4-(4-methyl-3-pyridylamino)thiobenzophenone,
 2,4'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)thiobenzophenone,
 2-Chloro-4'-fluoro-4-(4-isoquinolylamino)-2'-methyl(thiobenzophenone),
 4'-*n*-Butyl-2-chloro-4-(4-isoquinolylamino)-2'-methyl(thiobenzophenone),
 20 2-Chloro-4-(5-isoquinolylamino)-2'-methyl(thiobenzophenone),
 2-Chloro-4-(4-isoquinolylamino)-4'-methoxy-2'-methyl(thiobenzophenone),
 2-Fluoro-4-(4-isoquinolylamino)-4'-methoxy-2'-methyl(thiobenzophenone),
 2,4'-Dichloro-2'-methyl-4-(1-methyl-7-indolylamino)thiobenzophenone,
 2-Chloro-4-(1-methyl-7-indolylamino)-2',4',5'-trimethyl(thiobenzophenone),
 25 2-Chloro-2',5'-dimethyl-4-(1-methyl-7-indolylamino)thiobenzophenone,
 2-Chloro-4-(3-ethoxy-4-isoquinolylamino)-2'-methyl(thiobenzophenone),
 2-Chloro-4-(1-ethoxy-4-isoquinolylamino)-2'-methyl(thiobenzophenone),
 4-(2-Benzoxazolylamino)-2-chloro-2'-methyl(thiobenzophenone),
 4-(1-Methyl-2-benzimidazolylamino)-2-chloro-2'-methyl(thiobenzophenone),
 30 4-(2-Benzothiazolylamino)-2-chloro-2'-methyl(thiobenzophenone),
 2-Chloro-2'-methyl-4-(2-pyrimidylamino)thiobenzophenone,
 2-Chloro-2'-methyl-4-(7-methyl-purin-6-ylamino)thiobenzophenone,
 2-Chloro-2'-methyl-4-(2-methyl-5-benzothiazolylamino)thiobenzophenone,
 2-Chloro-2'-methyl-4-(pyrazin-2-ylamino)thiobenzophenone,
 35 2-Chloro-2'-methyl-4-(5-pyrimidylamino)thiobenzophenone
 and salts thereof with pharmaceutically acceptable acids, hydrates and solvates.

28. A compound of formula Ib selected from the group consisting of:
- 2-Chloro-2'-methyl-4-(4-pyridylamino)benzophenone oxime
 - 2-Chloro-2'-methyl-4-(2-pyridylamino)benzophenone oxime
 - 2-Chloro-2'-methyl-4-(5-nitro-2-pyridylamino)benzophenone oxime
 - 5 4-(6-Amino-5-nitro-2-pyridylamino)-2-chloro-2'-methylbenzophenone oxime
 - 6-Chloro-2-((3-chloro-4-((hydroxyimino)(2-methylphenyl)methyl))phenylamino)isonicotinic acid
 - 4-(6-carbonitrile-2-pyridylamino)-2-chloro-2'-methylbenzophenone oxime
 - 2-Chloro-2'-methyl-4-(3-pyridylamino)benzophenone oxime
 - 10 2-Chloro-4-(5,6-diamino-2-pyridylamino)-2'-methylbenzophenone oxime
 - 2-Chloro-2'-methyl-4-(3-nitro-2-pyridylamino)benzophenone oxime
 - 4-(3-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone oxime
 - 4-(5-Amino-2-pyridylamino)-2-chloro-2'-methylbenzophenone oxime
 - 2-Chloro-4-(4-isoquinolylamino)-2'-methylbenzophenone oxime
 - 15 t-Butyl 5-((3-chloro-4-((hydroxyimino)(2-methylphenyl)methyl))phenylamino) nicotinoate
 - 2-Chloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone oxime
 - 2-Chloro-2'-methyl-4-(6-methyl-3-pyridylamino)benzophenone oxime
 - 4-(5-Bromo-3-pyridylamino)-2-chloro-2'-methylbenzophenone oxime
 - 4-(5-Carbonitrile-3-pyridylamino)-2-chloro-2'-methylbenzophenone oxime
 - 20 4-(3-Bromo-2-pyridylamino)-2-chloro-2'-methylbenzophenone oxime
 - 2-Chloro-2'-methyl-4-(8-quinolylamino)benzophenone oxime
 - 2-Chloro-4-(6-ethoxy-3-pyridylamino)-2'-methylbenzophenone oxime
 - 4-(4-Bromo-1-isoquinolylamino)-2-chloro-2'-methylbenzophenone oxime
 - 2-Chloro-2'-methyl-4-(2-methyl-5-trifluoromethyl-3-pyridylamino)benzophenone oxime
 - 25 2-Chloro-2'-methyl-4-(3-quinolylamino)benzophenone oxime
 - 2-Chloro-4-(4-ethoxy-3-pyridylamino)-2'-methylbenzophenone oxime
 - 2-Chloro-4-(2-ethoxy-3-pyridylamino)-2'-methylbenzophenone oxime
 - 2-Chloro-2'-methyl-4-(2-methyl-6-quinolylamino)benzophenone oxime
 - 2-Chloro-2'-methyl-4-(2-quinolylamino)benzophenone oxime
 - 30 2-Chloro-4-(7-chloro-4-quinolylamino)-2'-methylbenzophenone oxime
 - 2-Chloro-2'-methyl-4-(4-quinolylamino)benzophenone oxime
 - 2-Chloro-2'-methyl-4-(1-methyl-7-indolylamino)benzophenone oxime
 - 2-Chloro-2'-methyl-4-(1-methyl-5-indolylamino)benzophenone oxime
 - 2-Chloro-2',5'-dimethyl-4-(4-methyl-3-pyridylamino)benzophenone oxime,
 - 35 2-Chloro-2',5'-dimethyl-4-(4-isoquinolylamino)benzophenone oxime,
 - 2-Chloro-4-(4-isoquinolylamino)-2',4',5'-trimethylbenzophenone oxime,
 - 2,3'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone oxime,
 - 2-Fluoro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone oxime,

- 2,4'-Dichloro-2'-methyl-4-(4-methyl-3-pyridylamino)benzophenone oxime,
 2-Chloro-4'-fluoro-4-(4-isoquinolylamino)-2'-methylbenzophenone oxime,
 4'-*n*-Butyl-2-chloro-4-(4-isoquinolylamino)-2'-methylbenzophenone oxime,
 2-Chloro-4-(5-isoquinolylamino)-2'-methylbenzophenone oxime,
 5 2-Chloro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone oxime,
 2-Fluoro-4-(4-isoquinolylamino)-4'-methoxy-2'-methylbenzophenone oxime,
 2,4'-Dichloro-2'-methyl-4-(1-methyl-7-indolylamino)benzophenone oxime,
 2-Chloro-4-(1-methyl-7-indolylamino)-2',4',5'-trimethylbenzophenone oxime,
 2-Chloro-2',5'-dimethyl-4-(1-methyl-7-indolylamino)benzophenone oxime,
 10 2-Chloro-4-(3-ethoxy-4-isoquinolylamino)-2'-methylbenzophenone oxime,
 2-Chloro-4-(1-ethoxy-4-isoquinolylamino)-2'-methylbenzophenone oxime,
 4-(2-Benzoxazolylamino)-2-chloro-2'-methylbenzophenone oxime,
 4-(1-Methyl-2-benzimidazolylamino)-2-chloro-2'-methylbenzophenone oxime,
 4-(2-Benzothiazolylamino)-2-chloro-2'-methylbenzophenone oxime,
 15 2-Chloro-2'-methyl-4-(2-pyrimidylamino)benzophenone oxime,
 2-Chloro-2'-methyl-4-(7-methyl-purin-6-ylamino)benzophenone oxime,
 2-Chloro-2'-methyl-4-(2-methyl-5-benzothiazolylamino)benzophenone oxime,
 2-Chloro-2'-methyl-4-(pyrazin-2-ylamino)benzophenone oxime,
 2-Chloro-2'-methyl-4-(5-pyrimidylamino)benzophenone oxime,
 20 and salts thereof with pharmaceutically acceptable acids, hydrates and solvates.
29. A pharmaceutical composition containing as an active component a compound according to any one of claims 1-28 together with a pharmaceutically acceptable excipient or carrier.
- 25 30. A composition according to claim 29 further comprising a second active ingredient optionally selected from the group consisting of glucocorticoids, vitamin D or vitamin D analogues, anti-histamines, platelet activating factor (PAF) antagonists, anticholinergic agents, methyl xanthines, β -adrenergic agents, salicylates, indomethacin, flufenamate, naproxen, timegadine, gold salts, penicillamine, serum cholesterol-reducing agents,
 30 retinoids, zinc salts, and salicylazosulfapyridin.
31. A composition according to claim 29 or 30 in unit dosage form containing between 0.05 and 1000 mg of the active ingredient.
- 35 32. Use of a compound according to any one of claims 1-28 for the preparation of a medicament for the treatment or prophylaxis of inflammatory diseases or conditions.

33. The use of claim 32, wherein the disease or condition is asthma, allergy, arthritis, including rheumatoid arthritis, osteoarthritis and spondyloarthritis, gout, atherosclerosis, inflammatory bowel disease, Crohn's disease, proliferative and inflammatory skin disorders, such as psoriasis, atopic dermatitis and acne vulgaris, uveitis, sepsis, septic shock, AIDS related diseases and osteoporosis.
34. A method for the treatment or prophylaxis of inflammatory diseases or conditions, the method comprising administering, to a patient in need thereof, an effective amount of one or more compounds according to any one of claims 1-28.
35. The method of claim 34 further comprising administering a second active component selected from the group consisting of glucocorticoids, vitamin D's, anti-histamines, platelet activating factor (PAF) antagonists, anticholinergic agents, methyl xanthines, β -adrenergic agents, salicylates, indomethacin, flufenamate, naproxen, tilmegadine, gold salts, penicillamine, serum cholesterol-reducing agents, retinoids, zinc salts, and salicylazosulfapyridin.
36. The method of claim 34, wherein the disease or condition is selected from the group consisting of asthma and allergy.
37. The method of claim 34, wherein the disease or condition is selected from the group consisting of arthritis, including rheumatoid arthritis, osteoarthritis and spondyloarthritis, gout, atherosclerosis, inflammatory bowel disease, Crohn's disease, uveitis, sepsis and septic shock.
38. The method of claim 34, wherein the disease or condition is a proliferative and/or inflammatory skin disorder such as psoriasis, atopic dermatitis and acne vulgaris.
39. A method for the treatment and/or prophylaxis of osteoporosis, the method comprising administering to a patient in need thereof an effective amount of one or more compounds according to any one of claims 1-28.
40. A method for the treatment of AIDS related diseases, the method comprising administering to a patient in need thereof an effective amount of one or more compounds according to any one of claims 1-28.
41. The method of any one of claims 34-40 comprising administering to a patient in need of systemic treatment a dose of from 0.01 to 400 mg/kg body weight, preferably a dose of from 0.1 to 100 mg/kg body weight one or more times daily of a compound according to any one of claims 1-28.